

Appendix E

Harris Environmental Group, Inc.
Draft Biological Assessment
TEP Proposed Sahuarita-Nogales
Transmission Line Project
Central Corridor (HEG 2003b)

BIOLOGICAL ASSESSMENT
OF THE
TUCSON ELECTRIC POWER
SAHUARITA – NOGALES TRANSMISSION LINE
CENTRAL CORRIDOR

DRAFT

15 MAY 2003

PREPARED FOR:

TUCSON ELECTRIC POWER

ONE SOUTH CHURCH

PO BOX 711

TUCSON, ARIZONA 85702

PREPARED BY:

HARRIS ENVIRONMENTAL GROUP

58 EAST 5TH STREET

TUCSON, ARIZONA 85705

(520) 628 – 7648

TABLE OF CONTENTS

EXECUTIVE SUMMARY.....	3
1.0 PROJECT DESCRIPTION.....	5
1.1 Proposed Action	5
1.3 Project Area	9
1.4 Conservation Measures.....	11
2.0 FEDERALLY LISTED SPECIES.....	15
2.1 Cactus ferruginous pygmy-owl (<i>Glaucidium brasilianum cactorum</i>) (Endangered)	17
2.1a Action Area	17
2.1b Natural History and Distribution:.....	17
2.1c Critical Habitat	19
2.1d Current Status Statewide	19
2.1e Environmental Baseline	20
2.1f Effects of Proposed Action on the CFPO.....	21
2.1g Cumulative Effects.....	24
2.1h Effects Determination and Incidental Take.....	24
2.2 Southwestern willow flycatcher (<i>Empidonax traillii extimus</i>) (Endangered)	25
2.2a Action Area	25
2.2b Natural History and Distribution.....	25
2.2c Critical Habitat	26
2.2d Current Status Statewide	26
2.2e Environmental Baseline	27
2.2f Effects of Proposed Action on the SWFL.....	28
2.2g Cumulative Effects.....	29
2.2h Effects Determination and Incidental Take.....	30
2.3 Lesser long-nosed bat (<i>Leptonycteris curasoae yerbabuenae</i>) (Endangered).....	31
2.3a Action Area	31
2.3b Natural History and Distribution.....	31
2.3c Critical Habitat	33
2.3d Current Status Statewide	33
2.3e Environmental Baseline	34
2.3f Effects of Proposed Action on LLNB.....	34
2.3g Cumulative Effects.....	36
2.3h Incidental Take.....	37
2.4 Pima pineapple cactus (<i>Coryphantha scheeri</i> var. <i>robustispina</i>) (Endangered)	38
2.4a Action Area	38
2.4b Natural History and Distribution.....	38
2.4c Critical Habitat	38
2.4d Current Status Statewide	38
2.4e Environmental Baseline	39
2.4f Effects of Proposed Action on the PPC	40
2.4g Cumulative Effects.....	41
2.4h Effects Determination	42
2.5 Jaguar (<i>Panthera onca</i>) (Endangered).....	43

2.5a	Action Area	43
2.5b	Natural History and Distribution.....	43
2.5c	Critical Habitat	44
2.5d	Current Status Statewide	44
2.5e	Environmental Baseline	44
2.5f	Effects of Proposed Action on the Jaguar	45
2.5g	Cumulative Effects	47
2.5h	Effects Determination and Incidental Take.....	47
2.6	Gila Topminnow (<i>Poeciliopsis occidentalis occidentalis</i>) (Endangered)	48
2.6a	Action Area	48
2.6b	Natural History and Distribution.....	48
2.6c	Critical Habitat	50
2.6d	Current Status Statewide	50
2.6e	Environmental Baseline	51
2.6f	Effects of Proposed Action on the Gila topminnow	52
2.6g	Cumulative Effects	52
2.6h	Effects Determination and Incidental Take.....	53
2.7	Mexican Gray Wolf (<i>Canis lupus baileyi</i>) (Endangered).....	54
2.7a.	Action Area	54
2.7b.	Natural History and Distribution	54
2.7c	Critical Habitat	55
2.7d	Current Status Statewide	55
2.7e	Environmental Baseline	55
2.7f	Effects of Proposed Action on the Mexican Gray Wolf	56
2.7g	Cumulative Effects	57
2.7h	Effects Determination and Incidental Take.....	57
3.0	USFS SENSITIVE SPECIES	58
3.1	Plants	64
3.2	Invertebrates	77
3.3	Birds.....	77
3.4	Reptiles and Amphibians.....	79
3.5	Mammals	81
4.0	BLM SENSITIVE SPECIES	83
4.1	Plants	84
4.2	Birds.....	85
4.3	Reptiles and Amphibians.....	86
4.4	Mammals	86
5.0	AGFD WILDLIFE OF SPECIAL CONCERN	89
6.0	LITERATURE CITED	96
7.0	LIST OF ACRONYMS	116
APPENDIX A	118
APPENDIX B	119
APPENDIX C	125
APPENDIX D	126

EXECUTIVE SUMMARY

Tucson Electric Power (TEP) and Citizens Communications (Citizens) are proposing to build a new, dual-circuit, 345,000-volt (345-kV) transmission line from the TEP South Substation in the vicinity of Sahuarita, Arizona to interconnect with Citizens system at a Gateway Substation that TEP will construct west of Nogales, Arizona. From the Gateway Substation, the proposed transmission line will continue south across the United States-Mexico border for approximately 60 miles (mi) (98 kilometers [km]) into the Sonoran region of Mexico, connecting with the Comisión Federal de Electricidad (CFE, the national electric utility of Mexico) at the Santa Ana Substation. The proposed transmission line will improve Citizens' service in Nogales and allow for the transfer of blocks of electrical energy between the United States and Mexico. Southern Arizona and Sonora, Mexico have experienced rapid growth, and forecasts predict this growth will continue. Citizens' customers have already experienced outages due to limited transmission facilities into the region. TEP recognizes the need to improve transmission into the southern Arizona region and proposes to assist Citizens in meeting an Arizona Corporation Commission (ACC) mandate to improve the reliability and service of its Nogales electrical system. The ACC has ordered Citizens to improve its system by the end of 2003. The TEP Sahuarita – Nogales Transmission Line, a double-circuit 345-kV transmission line will provide the additional reliability that Citizens requires while providing additional capacity into the southern Arizona region for future needs.

This Biological Assessment (BA) was prepared to meet the requirements of Section 7 of the Endangered Species Act (ESA) of 1973, 16 U.S.C. Section 1536(a)(2). Section 7 requires all federal agencies to consult with the United States Fish and Wildlife Service (USFWS) if an action may affect listed species or their designated critical habitat. Section 7 consultation is required for any project that requires a federal permit or receives federal funding. Action is defined broadly to include funding, permitting, and other regulatory actions. All activities associated with construction of the TEP Sahuarita – Nogales Transmission Line are included in the proposed action being evaluated for this BA. Because TEP has applied for a Presidential Permit to construct the transmission line across the international border, the Department of Energy (DOE) is preparing a Draft Environmental Impact Statement (DEIS) (Tetra Tech 2003) concurrently with this document.

Federal agencies must ensure that any action they authorize, fund, or carry out is not likely to jeopardize the continued existence of a listed species, or result in the destruction or adverse modification of designated critical habitat. This is accomplished through consultation with the USFWS. If such species may be present, the applicant must conduct a BA to determine if a proposed action is likely to adversely affect listed species or designated critical habitat. USFWS will review this BA and issue a biological opinion (BO). DOE is the permitting agency for this proposed action, and therefore the lead federal agency in Section 7 consultation with USFWS.

The proposed action crosses a variety of land jurisdictions: including private, Arizona State Land Department (ASLD), Bureau of Land Management (BLM), and United States Department of Agriculture Forest Service (USFS). Because each jurisdiction has different requirements for environmental review of the proposed action, this document is subdivided by agency. SECTION 2 addresses species that receive protection under the ESA of 1973. SECTION 3 reviews the potential effects of the proposed action on those species classified as “Sensitive” by the USFS. SECTION 4 reviews the potential effects of the proposed action on those species classified as “Sensitive” by the BLM. SECTION 5 addresses those species that are considered “Wildlife of Special Concern” by the Arizona Game and Fish Department (AGFD). Because habitats often overlap different jurisdictions, many species have classifications within each agency. In these instances, the species is evaluated under the jurisdiction which affords the highest level of protection.

We contacted federal (USFWS) and state (AGFD) natural resource agencies to request information on possible special status species (sensitive, threatened, and endangered) that may exist on or near the proposed Central Corridor of the TEP Sahuarita – Nogales Transmission Line from Sahuarita to Nogales, Arizona. Agency correspondence is presented in Appendix A.

SUMMARY OF DETERMINATIONS FOR FEDERALLY LISTED SPECIES

Based on contact with USFWS, USFS, BLM, and AGFD, 7 federally listed species may be affected by the proposed action. Upon review of the current status of these species, the environmental baseline of the project area, the effects of the proposed actions on the species as well as cumulative effects, the following determinations are made for the 7 affected species (Table 1).

Table 1. Effects of the proposed action on federally listed species.

SPECIES	POTENTIAL EFFECT
<i>Cactus ferruginous pygmy-owl</i>	The proposed action may affect and is likely to adversely affect this species.
<i>Southwestern willow flycatcher</i>	The proposed action may affect but is not likely to adversely affect this species.
<i>Lesser long-nosed bat</i>	The proposed action may affect and is likely to adversely affect this species.
<i>Pima pineapple cactus</i>	The proposed action may affect and is likely to adversely affect this species.
<i>Jaguar</i>	The proposed action may affect but is not likely to adversely affect this species.
<i>Gila topminnow</i>	The proposed action may affect but is not likely to adversely affect this species.
<i>Mexican gray wolf</i>	The proposed action may affect but is not likely to adversely affect this species.

1.0 PROJECT DESCRIPTION

1.1 PROPOSED ACTION

The proposed TEP Central Corridor Sahuarita – Nogales Transmission Line will consist of 12 transmission line wires, or conductors, and 2 neutral ground wires that will provide lightning protection and fiber optic communication, on a single set of support structures. The transmission line will originate at the existing South Substation, in the vicinity of Sahuarita, Arizona, and interconnect with Citizens system at a Gateway Substation that TEP will construct west of Nogales, Arizona. The double-circuit transmission line will continue from the Gateway Substation south to cross the United States-Mexico border and extend approximately 60 mi (98 km) into the Sonoran region of Mexico, connecting with the Comisión Federal de Electricidad (CFE, the national electric utility of Mexico) at the Santa Ana Substation. Figure 1 shows the overall proposed project location.

The South Substation in Sahuarita will be upgraded and expanded to provide interconnection between a new TEP 345-kV transmission line and the new Gateway Substation west of Nogales. The South Substation will be expanded by approximately 1.3 acres (0.53 ha) to add a switching device that will connect to the proposed transmission line, with a 100 ft (30 m) expansion of the existing fence line for the addition of the second 345-kV circuit. The new Gateway Substation will include a 345-kV to 115-kV power transformer to provide power to the local area. The new Gateway Substation will be constructed within a developed industrial park north of Mariposa Road (State Route 189), approximately 0.5 mi (0.8 km) east of the Coronado National Forest (CNF) boundary (Northeast $\frac{1}{4}$ of Section 12, Township 24 South, Range 13 East). The TEP portion of the site is approximately 18 acres (7.3 ha) and is within the City of Nogales, Arizona. TEP has purchased the substation site and preliminary construction activities have been completed. TEP is flexible in the placement of a fiber-optic regeneration site, but it will likely be located in the area of Township 18 South, Range 12 East, approximately 10 mi (16 km) southwest of Sahuarita on private land. The fiber optic regeneration site will consist of an approximate 0.5-acre (0.2-ha) fenced yard, containing a 10 ft (3 m) by 20 ft (6 m) concrete pad with an equipment house. The cleared area for the equipment house will be approximately 20 ft (6 m) by 30 ft (9 m). There will be three 3-acre (1.2-ha) construction staging areas (located near the South and Gateway Substations and the Interstate 19 [I-19]/Arivaca Road interchange) and an 80 acre (32 ha) temporary laydown yard (also near the I-19/Arivaca Road interchange) used during construction of the proposed line.




Figure 1. Map of the TEP Sahuarita – Nogales Transmission Line Central Corridor.

The primary support structures to be used for the transmission line are self-weathering steel single structures, or monostructures (Figure 2). Dulled, galvanized steel lattice towers (Figure 3) will be used in locations where their use will minimize overall environmental impacts, in accordance with Arizona Corporation Commission (ACC) Decision No. 64356 (ACC 2001).





Figure 2. Monopole Transmission Line Structure Drawing and Photo.




Figure 3. Lattice Tower Transmission Line Structure Drawing and Photo.

1.2 PROJECT LOCATION

The Central Corridor extends for approximately 57.1 mi (91.9 km), from the South Substation to the United States-Mexico border, including 43.2 mi (69.5 km) along the El Paso Natural Gas Company (EPNG) gasline right-of-way (ROW). The length of the Central Corridor is approximately 15.1 mi (24.3 km) within the CNF, and approximately 1.25 mi (2.01 km) on BLM land. The Central Corridor will require approximately 390 support structures, including approximately 102 within the CNF and 9 on BLM land. The Central Corridor exits the TEP South Substation located within the incorporated area of the Town of Sahuarita and proceeds westerly for approximately 1.0 mi (1.6 km) before turning south for 1.5 mi (2.4 km). The corridor turns west across I-19 and continues through Pima County to the southwest, crossing approximately 1.25 mi (2.01 km) of federal land managed by BLM parallel to 2 existing TEP transmission lines (138-kV and 345-kV). The Central Corridor turns south to parallel the EPNG gasline ROW until reaching approximately 3 mi (4.8 km) south of the existing TEP Cyprus Sierrita Substation. South of the TEP Cyprus Sierrita Substation, the Central Corridor continues south to follow the EPNG gasline ROW to the south.

The Central Corridor continues south about 1.0 mi (1.6 km) west of I-19, and passes Amado, Tubac, and Tumacacori. The Central Corridor continues approximately 2.0 mi (3.2 km) south of Tumacacori, and enters the CNF, adjacent to the EPNG gasline ROW. The Central Corridor centerline diverges from the EPNG gasline ROW for approximately 1.9 mi (3.1 km) and avoids the USFS inventoried roadless area (IRA). The Central Corridor continues through the CNF, paralleling the EPNG gasline ROW to the southeast for several miles to the forest boundary. The proposed corridor exits CNF onto private land and proceeds 0.5 mi (0.8 km) east to the Gateway Substation. From the Gateway Substation, the proposed corridor returns to the west through private land and then turns south to parallel the CNF boundary. The proposed corridor meets the United States-Mexico border approximately 3,300 ft (1,006 m) west of Arizona State Highway 189 in Nogales, Arizona.

TEP will use existing access when feasible. Approximately 13.8 mi (22.2 km) of temporary new roads will be built for construction of the Central Corridor on the CNF (URS 2003a); spur roads off existing access roads adjacent to TEP transmission lines will provide project access on BLM land. Transmission line tensioning and pulling and fiber-optic splicing sites will also disturb land. The total new temporary area of disturbance on the CNF during construction of the Central Corridor will be approximately 105 acres (42.5 ha) (URS 2003a). Following construction, TEP will close new roads, construction areas, and existing roads not required for project maintenance in accordance with agreements with land owners or managers (e.g., BLM or USFS). On national forest land, TEP will close existing road mileage equal to that required for project maintenance, to avoid impacting the current road density. The maintenance access required by TEP will be limited to roads to selected structures, rather than a single cleared ROW leading to the United States-Mexico border. On the CNF transmission line tensioning and pulling sites, fiber-optic splicing sites, and construction yard areas will be obliterated within six months of the project becoming fully operational (URS 2003a).

1.3 PROJECT AREA

The project area includes the location where all construction and associated activities will occur along the ROW. Action areas are locations affected directly or indirectly by these activities and often include sites outside the immediate area of construction. Action areas are unique for each listed species and are outlined in SECTION 2.0 of this document.

Between Sahuarita and Nogales, the proposed action crosses four distinct biotic communities, or biomes (Brown 1994). A complete list of plant species documented during field surveys in 2002 is presented in Appendix B.



Figure 4. Sonoran desertscrub.

The northern end of the corridor contains vegetation characteristic of the Sonoran desertscrub biome (Figure 4). This biome is typically represented by saguaro (*Carnegiea gigantea*), cholla and prickly pear (*Opuntia* spp.) cacti, ocotillo (*Fouquieria splendens*), mesquite, (*Prosopis velutina*), acacia (*Acacia* spp.) paloverde (*Parkinsonia* spp.), creosote (*Larrea tridentata*), triangle-leaf bursage (*Ambrosia deltoidea*), and brittlebush (*Encelia farinosa*).

Vegetation south of the ASARCO mine transitions into the semidesert grassland biome (Figure 5). This area is dominated by grama (*Bouteloua* spp.), lovegrass (*Eragrostis* spp.), and three-awn (*Aristida* spp.) grasses, with low shrubs such as mesquite and acacia locally co-dominant. Agave (*Agave* spp.) and yucca (*Yucca* spp.) are also common in this biome. These grasslands are transected by desert riparian scrub dominated by mesquite and netleaf hackberry (*Celtis reticulata*).



Figure 5. Semidesert grassland.



Figure 6. Madrean oak woodland.

The higher elevations (above 3,500 ft [1,067 m]) of the project area are within the madrean oak woodland biome (Figure 6). Representative plants of this biome within the project area include Mexican blue oak (*Quercus oblongifolia*) and emory oak (*Q. emoryi*) trees, side-oats grama (*B. curtipendula*), hairy grama (*B. hirsuta*), and fluffgrass (*Erioneuron pulchellum*).

The 4th biome represented within the project area is the Sonoran deciduous riparian forest (Figure 7), which is located within or near the ROW in Peck Canyon and in the Santa

Cruz River. The high water table in these areas supports stands of Fremont cottonwood (*Populus fremontii*), ash (*Fraxinus pennsylvanica* ssp. *velutina*), walnut (*Juglans major*), and willow (*Salix* spp.) trees.

The proposed ROW begins at an elevation of approximately 2,674 ft (815 m) at the TEP South Substation and reaches its maximum elevation of approximately 4,321 ft (1,317 m) near Tinaja Peak located southwest of the ASARCO Mine complex.

The Chiltepine Botanical Area is a 2,836 ac (1,148 ha) reserve located approximately 2 mi (1.2 km) west of the Central Corridor, in the northern portion of the Tumacacori Ecosystem Management Area (EMA) of the CNF. This area was established in June 1999 for the purpose of protecting and facilitating the study of chiltepine. These wild chiles typically are found in the more tropical environments between Mexico and South America. This area has been noted as the northernmost occurrence of chiltepine in the world.



Figure 7. Sonoran deciduous riparian forest.

Between 12 June and 22 June 2002, the Walker Fire, a human-caused fire, burned 16,369 ac (6,624 ha) of land along the United States-Mexico border approximately 1mi (1.6 km) west of the southern end of the Central Corridor. Portions of the Walker fire were very hot, especially near the international border and the upper slopes of ridges, while other areas, like Walker Canyon, burned relatively cool (T. Newman, CNF, pers. comm., 26 November 2002). While vegetation has begun to recover in some areas, other areas are highly susceptible to erosion due to reduced groundcover (Figure 8).



Figure 8. Area burned in Walker fire.

1.4 CONSERVATION MEASURES

PROJECT-WIDE CONSERVATION MEASURES

1. Environmental Training - All construction supervisors will be required to attend environmental training, which will outline their obligation to obey applicable laws and regulations regarding wildlife and habitats (Appendix C).
2. Erosion Control Measures - TEP is in consultation with CNF regarding development of Best Management Practices (BMPs) for minimizing proposed project impacts on geologic, soil, and water resources on national forest land, in accordance with the USFS "Soil and Water Conservation Practices Handbook" (USFS 1990). Specific BMPs will be identified after coordination with Arizona Department of Environmental Quality (ADEQ) and before implementation of the project, for the entire length of the selected corridor.
3. Fire Prevention Plan - A Fire Prevention Plan is under development to minimize the risk of accidental wildfire. All construction activities will adhere to this plan and fire suppression equipment will be available to all work crews. On CNF lands, the Fire Prevention Plan will comply with Forest Service Manual 5100.
4. Hazardous Material Spill Response Plan - A Hazardous Material Spill Response Plan is under development which will describe the measures and practices to prevent, control, cleanup, and report spills of fuels, lubricants, and other hazardous substances during construction operations. This plan will ensure that no hazardous materials are stored, dispensed, or transferred in streams, watercourses, or dry washes, and vehicles are regularly inspected and maintained to prevent leaks.
5. Invasive Species Control - An Invasive Species Management Plan in accordance with Executive Order 13112 is under development in coordination with CNF, ASLD, and BLM to identify problem areas and mitigation measures.
6. Road Closure/Obliteration - TEP has committed to obliterate and permanently close 1 mi (1.6 km) of existing road on CNF (to be identified by CNF) for every 1 mi (1.6 km) of proposed new road used in the construction, operation, or long-term maintenance of the proposed action. TEP will monitor road closures during regularly scheduled inspection flights and/or ground inspections, and repair or replace road-closure structures as necessary following construction. Furthermore, TEP will cooperate with landowners on all ongoing road closure maintenance.

The following selective criteria and techniques for closing roads are taken from Section 1.3.2 of the Roads Analysis (URS 2003) and applies to access roads on CNF. Administrative roads will be closed to the general public but made available to TEP and its assigned contractors for the evaluation, maintenance, or upgrading of existing facilities.

Closure methods for administrative roads will include the following:

- a. Placement of heavy pipe posts with an attached, locked chain entrance on the road.
- b. Placement of heavy pipe posts with an attached, locked gate in a manner that blocks entrance on the road.
- c. Placement of a pipe barricade across the roadbed, locked in place in multiple locations in concrete sleeves.

The following methods may be used for the long-term closure of transmission line access roads used during construction and those roads required to be closed by the CNF. These roads may be reopened for emergency repair of transmission facilities, but will not be used intermittently as with administrative roads. Techniques include:

- a. Placement of boulders or other natural impediments across the road.
- b. Placement of a berm or trench across the the road.
- c. Rip, obliterate, and reseed/revegetate portions of roadbed as needed. This effort could be applied to the initial visual portion of roadway (e.g., first 100 ft [30 m]) to effectively obscure the roadway. This could be accomplished by transplanting native species of medium and large vegetation from the general area and reseeding with native grasses. By obscuring visible portions of roadway, future vehicular travel could be more effectively discouraged than by placing berms or other unnatural impediments to an otherwise visually inviting roadway.

7. Additional mitigation measures are outlined in Table 2.2-2 of the DEIS (Tetra Tech 2003).

SPECIES-SPECIFIC CONSERVATION MEASURES

Cactus ferruginous pygmy-owl (CFPO)

1. Protocol surveys – 2 consecutive years of protocol surveys must be conducted before construction activities can begin within 1,312 ft (400 m) of designated habitat. If a CFPO is detected, USFWS has determined that certain continued construction activities will not harm or harass a CFPO as defined by ESA regulations. In areas where two consecutive years of protocol surveys cannot be completed, construction will occur outside of the breeding season.

Four zones are described (Zone I through Zone IV) that are based upon the distance of construction activity from a known nest or activity center. Certain levels of construction can occur within each zone without resulting in harm or harassment of the species. Situations that do not comply with the restrictions provided for each zone will require USFWS authorization before construction continues. Specific

development restrictions that apply to each of the four zones are described in the sections below:

Zone I: 0 to 328 ft (100 m) from the CFPO Activity Center

1. No additional clearing of vegetation will be permitted without authorization from USFWS and relevant land management agencies.
2. Construction-related activities may continue on land that has been cleared of vegetation provided that they do not exceed the level and/or intensity of activity that was occurring during the period of time that the territory was established.
3. Activities that will be more intense or cause more noise disturbance than was occurring during the period of time that the territory was established cannot proceed without authorization from USFWS and relevant land management agencies.

Zone II: 328 ft (100 m) to 1,312 ft(400 m) from the CFPO Activity Center

1. No additional clearing of vegetation will be permitted without authorization from USFWS and relevant land management agencies.
2. No restrictions on the nature or type of construction activity (excluding the clearing of vegetation) from 1 August through 31 January of the following calendar year.
3. Construction activities during the breeding season (1 February to 31 July) cannot exceed the levels or intensity of activities that occurred at the time the territory was established.

Zone III: 1,312 ft (400 m) to 1,969 ft (600 m) from the CFPO Activity Center

1. No additional clearing of vegetation will be permitted without authorization from USFWS and relevant land management agencies.
2. No restrictions on the levels or intensity of construction activity (excluding the clearing of vegetation) at any time of the year.

Zone IV: Greater than 1,969 ft (600 m) from the CFPO Activity Center

1. No restrictions – any activity consistent with the project description provided to USFWS (as amended by supplemental reports) is allowed. For the purposes of this consultation, USFWS assumes that all construction or construction-related activities referred to under each zone description will be limited to those described in the project description in this BA.
2. All saguaros within construction areas will be transplanted or mitigated with

minimum 6.5 ft (2 m) specimens. Within riparian desertscrub and deciduous riparian areas, tree and shrub removal will be minimized to the greatest extent possible.

Lesser long-nosed bat (LLNB)

1. Agave within construction areas will be transplanted or replaced with similar age and size class individuals.

Pima pineapple cactus (PPC)

1. Purchase of credits in a USFWS-approved conservation bank for PPC at a ratio to be determined in consultation with USFWS.

Jaguar

1. Two remote cameras will be donated to the Jaguar Conservation Team to assist with monitoring of jaguar movements across the Arizona-Mexico border. These cameras will all be placed within the Tumacacori EMA under permit from CNF. If female jaguar or cubs are documented by the Jaguar Management Team within the Tumacacori EMA, consultation with USFWS will be reinitiated.

2.0 FEDERALLY LISTED SPECIES

Special status species are plant and wildlife species that are of concern because their populations are either in jeopardy of extinction or are declining in number. The AGFD and USFWS were contacted concerning information on possible threatened and endangered species that may exist on or near the proposed action. In a letter dated 14 June 2002, USFWS listed 17 endangered species, seven threatened species, and two proposed species that occur in Pima and Santa Cruz counties, Arizona (Table 2). Agency correspondence is presented in Appendix A. Species included in USFWS correspondence, but excluded from evaluation are addressed in Appendix D.

Meetings with USFWS and USFS personnel were held on 9 April, 13 May, 3 December 2002, and 28 March 2003 to discuss the potential effects of the proposed action on special status species. BLM personnel also attended the 3 December 2002 meeting. Additional meetings were held with USFWS on 30 May, 6 November, 10 December 2002, and 19 March 2003, and with AGFD on 19 April 2002.

Table 2. Federally listed species that may occur near the proposed action.

SPECIES	STATUS	<i>DRAFT</i> DETERMINATION
Canelo Hills ladies' tresses	Endangered	No Effect
Cactus ferruginous pygmy-owl	Endangered	May affect, likely to adversely affect
Desert pupfish	Endangered	No Effect
Gila topminnow	Endangered	May affect, not likely to adversely affect
Huachuca water umbel	Endangered	No Effect
Jaguar	Endangered	May affect, not likely to adversely affect
Jaguarundi	Endangered	No Effect
Kearney's blue star	Endangered	No Effect
Lesser long-nosed bat	Endangered	May affect, likely to adversely affect
Masked bobwhite	Endangered	No Effect
Mexican gray wolf	Endangered	May affect, not likely to adversely affect
Nichols turk's head cactus	Endangered	No Effect
Northern aplomado falcon	Endangered	No Effect
Ocelot	Endangered	No Effect
Pima pineapple cactus	Endangered	May affect, likely to adversely affect
Sonoran pronghorn	Endangered	No Effect
Sonoran tiger salamander	Endangered	No Effect
Southwestern willow flycatcher	Endangered	May affect, not likely to adversely affect
Bald eagle	Threatened	No Effect
Brown pelican	Threatened	No Effect
Chiricahua leopard frog	Threatened	No Effect
Loach minnow	Threatened	No Effect
Mexican spotted owl	Threatened	No Effect
Sonora chub	Threatened	No Effect
Spikedace	Threatened	No Effect
Mountain plover	Proposed	No Effect
Gila chub	Proposed	No Effect

2.1 CACTUS FERRUGINOUS PYGMY-OWL (*Glaucidium brasilianum cactorum*) (Endangered)

2.1a Action Area

The action area includes all areas potentially affected, directly or indirectly, by all aspects of the project. The action area for the CFPO includes those areas of habitat below 4,000 ft (1,219 m) that may be directly impacted by construction as well as potential nesting sites within 1,312 ft (400 m) of the proposed action (USFWS 2000) that may be subject to noise disturbance during construction. In addition, an 7.08 mi (11.4 km) buffer area surrounding the project area is included in the action area because juvenile CFPO have been documented traveling up to 7.08 mi (11.4 km) during dispersal (M. Wrigley, USFWS, pers. comm., May 2001).

2.1b Natural History and Distribution:

USFWS listed CFPO in Arizona on 10 March 1997 (USFWS 1997a) as endangered. Listing was based on historical and current evidence that suggested a significant population decline of this subspecies had occurred in Arizona. USFWS considered the loss and alteration of habitat as the primary threat to the remaining population. A recovery plan for the species is currently in development by the CFPO recovery team.

CFPO (Figure 9) are small brown birds, with a cream-colored belly streaked with paler brown (Pyle 1997). The *cactorum* race; however, is described as “a well-marked, pale grayish extreme for the species” (Phillips et al. 1964). The call for this mostly diurnal owl is heard chiefly near dawn and dusk. The best field identification features are its small size, eyespots on the nape of the neck, and long reddish-barred tail, which is often nervously wagged or twitched (Monson 1998).

Originally CFPO were described as a separate subspecies based on specimens from Arizona and Sonora, Mexico. CFPO were first documented in the United States from a collection by Lieutenant Charles E. Bendire on 24 January 1872 in the “heavy mesquite thickets along Creek” near the present day site of historic Camp Lowell, Tucson (Coues 1872, Bendire 1892).



Figure 9. Cactus ferruginous pygmy-owl.

Very little is known about the life history of CFPO in Arizona (Cartron et al. 2000a). Little or no literature currently exists concerning life history variables such as longevity, age distribution, and recruitment. Current studies undertaken by AGFD, USFWS, and The University of Arizona are examining these variables.

The diet of CFPO is not well understood, but they are believed to be prey generalists (Cartron et al. 2000a). Observations, stomach content analysis, and records of Texas pygmy-owls suggest that these owls have a diverse diet that includes mammals, birds, reptiles, and insects (Proudfoot and Beasom 1997).

CFPO nest in cavities of larger trees (typically defined as a tree with a trunk at least 6 in [15 cm] diameter at breast height [DBH]) or large columnar cactus. Cavities may be naturally formed (e.g. knotholes) or excavated by woodpeckers. CFPO do not construct their own nest holes. All currently known CFPO nest sites in Arizona are in woodpecker excavated cavities in saguaros. Historically, the species also has been documented nesting in cottonwood, paloverde, and mesquite trees in Arizona.

Nesting activity for this owl species in Arizona begins in late winter to early spring (Lesh and Corman 1995, Abbate et al. 1996). Little is known about its courtship flight behavior. Egg laying begins by late April with three to four eggs typically laid. It is uncertain if only one brood is hatched per year. Nestlings have been observed through the end of July. During nesting, the male brings food to the female and young (Glinski 1998).

Historically, CFPO occurred from the lowlands of central Arizona, south through western Mexico to the states of Colima and Michoacan, and from southern Texas south through the Mexican states of Tamaulipas and Nuevo Leon. In Arizona, the species was documented as far north as New River and Cave Creek in northern Maricopa County (Harris and Duncan 1999). Elsewhere in Maricopa County, the species has been found near the Yuma County line along the Gila River at Agua Caliente, along the Salt River at Phoenix, and near the Verde River confluence. The eastern most verifiable record was along the Gila River at Old Fort Goodwin, located approximately 2 mi (1.2 km) southwest of present day Geronimo, Graham County, Arizona (Aiken 1937). In the southeastern part of the state, the species has been documented in recent times near Dudleyville along the lower San Pedro River between 1985 and 1987 (Harris and Duncan 1999), and probably also along lower Aravaipa Creek in 1987 (Monson 1987). Other localities in south central Arizona include historical records in Pinal County near Sacaton and Blackwater on the Gila River Indian Reservation, and at Casa Grande (Harris and Duncan 1999). Near the Mexican border, the species has been found in Santa Cruz County near Patagonia and in Sycamore Canyon west of Nogales. A likely accidental sighting was documented once on 10 April 1955 in eastern Yuma County near the Mexican border at Cabeza Prieta Tanks on the Cabeza Prieta National Wildlife Refuge (Monson and Phillips 1981, Harris and Duncan 1998).

Surveys conducted by University of Arizona biologists in Sonora, Mexico found 280 CFPO during the 2000 survey season. CFPO within Sonora, Mexico and Arizona may have been the same population prior to agricultural expansion within the last 75 years. However, due to isolation, the genetic connection of the Arizona population to owls in the nearby state of Sonora, Mexico may be tenuous (USFWS 2002a).

CFPO have been documented in several habitat types in the northern portion of its range in Arizona and adjacent Mexico. In Arizona, these include streamside Sonoran riparian deciduous forest and woodland associations and Sonoran desertscrub. CFPO also inhabit Sinaloan deciduous forest and thornscrub in Mexico (not discussed here). The streamside associations include such species as cottonwood, ash, netleaf hackberry, willows, velvet

mesquite, and others. The Sonoran desertscrub associations are composed of relatively dense saguaro cactus stands associated with short trees such as palo verde, mesquite, and ironwood (*Olneya tesota*), and an open understory of triangle-leaf bursage, creosote, and various other cacti and shrubs. Throughout its range, CFPO occur at low elevations, generally below 4,000 ft (1,219 m).

CFPO found in Sonoran desertscrub habitats are typically associated with structurally diverse stands of desert riparian scrub with saguaros along washes (Wilcox et al. 2000). Such habitat is often referred to as xeroriparian vegetation (Johnson and Haight 1985). These washes have no permanent water flow. Instead, flow is intermittent and based on seasonal rainfall as well as strength and duration of individual storms. Desert riparian scrub vegetation is easily recognizable by the presence of a linear assemblage of trees and shrubs that grow along the wash. Density is higher and taller than the sparse desertscrub vegetation that typically exists in the adjacent uplands. Before listing the species as endangered, all known CFPO were documented in such Sonoran desertscrub habitat (Lesh and Corman 1995, Abbate et al. 1996).

At the northern periphery of the subspecies range in southern Arizona, CFPO distribution and preferred habitat is not well understood. It is believed CFPO require the cover of denser wooded areas with understory thickets, like riparian habitat, for nesting, foraging, and predator avoidance (Abbate et al. 2000). Riparian habitat also is known for its high density and diversity of animal species that constitute the prey base of CFPO.

A significant decline in the Arizona population has occurred over the past several decades (USFWS 1997a, Richardson et al. 2000). Loss or modification of habitat from woodcutting, agriculture, groundwater pumping, and related human activities has presumably contributed to the population decline (USFWS 1997a).

2.1c Critical Habitat

On 12 July 1999, USFWS designated approximately 731,712 acres (296,113 ha) of critical habitat supporting riverine, riparian, and upland vegetation in seven critical habitat units, located in Pima, Cochise, Pinal, and Maricopa counties of Arizona (USFWS 1999). However, on 21 September 2001, the U.S. District Court for the State of Arizona vacated this final rule designating critical habitat for CFPO, and remanded its designation back to the USFWS for further consideration. On 27 November 2002, USFWS proposed designating 1.2 million acres (485,000 ha) of critical habitat for CFPO in southern Arizona (Federal Register Vol. 67, No 229:71031-71064). The proposed action does not enter any areas proposed as critical habitat.

2.1d Current Status Statewide

USFWS determined that CFPO in Arizona were endangered because of the following factors (USFWS 1997a):

- present or threatened destruction, modification, or curtailment of its habitat or range;
- inadequacy of existing regulatory mechanisms;

- other natural and manmade factors, which include low genetic viability.

Surveys conducted statewide during the 2002 season confirmed a total of 18 adult CFPO and three nests in Arizona. Similar to the previous four years, there was greater than 50 percent fledgling mortality documented in 2002, with only one juvenile confirmed surviving dispersal (S. Richardson, USFWS, pers. comm., 3 December 2002).

One of most urgent threats to CFPO in Arizona is thought to be the loss and fragmentation of habitat (USFWS 1997a, Abbate et al. 1999). The complete removal of vegetation and natural features required for many large-scale and high-density developments directly and indirectly impacts CFPO survival and recovery (Abbate et al. 1999). In recent decades, CFPO riparian habitat has continually been modified and destroyed by agricultural development, woodcutting, urban expansion, and general watershed degradation (Phillips et al. 1964, Brown et al. 1977, State of Arizona 1990, Bahre 1991, Stromberg et al. 1992, Stromberg 1993a and 1993b). Sonoran desertscrub has been affected to varying degrees by urban and agricultural development, woodcutting, and livestock grazing (Bahre 1991). Pumping of groundwater and the diversion and channelization of natural watercourses are also likely to have reduced CFPO habitat.

Proudfoot and Slack (2001) found that CFPO in northwestern Tucson may be isolated from other populations in Arizona and Mexico. Low genetic variability can lead to a reduction in reproductive success and environmental adaptability. In 1998 and 1999, two cases of sibling CFPO pairing and breeding were documented (Abbate et al. 1999). In both cases, young were fledged from the nesting attempts. These unusual pairings may have resulted from extremely low numbers of available mates within dispersal range, and/or from barriers (including fragmentation of habitat) that have influenced dispersal and limited the movement of young owls (Abbate et al. 1999).

Soule (1986) notes that very small populations are in extreme jeopardy due to their susceptibility to a variety of factors, including variations in birth and death rates that can result in extinction. In small populations such as with CFPO, each individual is important for its contribution to the genetic variability of that population.

2.1e Environmental Baseline

CFPO habitat north of Sahuarita Road consists of Sonoran desertscrub with relatively high species diversity and structural diversity, including scattered saguaro cacti containing potential nesting cavities. This area is within Survey Zone 1 (USFWS 2000) and has the highest potential for occupancy of the entire action area. Land status in this area is a mixture of private and state land. The Mission Mine Complex also is located within this section of the proposed action and grazing in the area.

CFPO habitat south of Sahuarita Road consists of Sonoran desertscrub dominated by mesquite and acacia trees, mixed with saguaro cacti, including non-native Lehmann's lovegrass (*Eragrostis lehmanniana*).



primarily undeveloped, but does contain some existing electrical distribution lines and associated roads (Figure 10) as well as low density housing developments. These grasslands are transected by desert riparian scrub dominated by mesquite and netleaf hackberry trees. Some areas of deciduous riparian forests are also found south of Arivaca Road in Sopori Wash and Peck Canyon. Land jurisdictions in this area include private, state, BLM, and USFS.

Figure 10. Example of existing disturbance within the corridor.

CFPO surveys were conducted by Harris Environmental Group, Inc. (HEG) biologists in 2001 and 2002 (data previously submitted to USFWS) in accordance with the approved protocol (USFWS 2000). Surveys were conducted in Sonoran desertscrub habitat where saguaros were present and in desert riparian scrub and deciduous riparian habitats that contained large trees (over 6 in [15.2 cm] DBH). No CFPO were detected during either survey year.

The only historical records of CFPO within the Nogales Ranger District (RD) of the CNF are in Sycamore Canyon (CNF 2000) and a dispersing juvenile in the Jarillas Alloment. USFS surveys in Sycamore Canyon in 1997 and 1998 did not locate CFPO. Additionally, USFS personnel surveyed 2,300 acres (930 ha) in 1999 with negative results and conducted 58 habitat assessments for CFPO habitat (CNF 2000). The habitat assessments identified four areas that ranked high enough to warrant CFPO surveys. No CFPO have been detected during surveys of these four areas (T. Newman, CNF, pers. comm., 9 October 2002).

2.1f Effects of Proposed Action on the CFPO

Direct Effects

Vehicle and Powerline Collisions

CFPO collisions with windows and fences have been documented in the Tucson area (USFWS 2002a), and observations of low flying CFPO across roadways indicate vehicle collisions are a realistic hazard (Abbate et al. 1999). While CFPO may be active during daylight, no CFPO have been detected within the action area, therefore, CFPO collisions with construction related vehicles are unlikely.

There is a small risk of a CFPO collision with power lines, however, raptors have lower rates of collision with power lines than passerine birds (McNeil et al. 1985). This reduced collision rate may be due to the visual acuity, maneuverability, and non-flocking tendencies (Nobel 1995). To minimize the risk of powerline collisions, TEP will construct the proposed transmission line following the guidelines outlined in "Suggested Practices for Raptor Protection on Power Lines: the State of the Art in 1996" (APLIC 1996).

Electrocution

Because power structures and towers are attractive perching and nesting sites for some raptor species, significant raptor mortality from electrocution has been reported in North

America (Harness and Wilson 2000). Electrocution occurs when a bird simultaneously touches two phase conductors or a conductor and a ground wire (Bevanger 1994). Most electrocutions occur on distribution lines (34-kV or less) rather than on transmission lines (69-kV or more), primarily because clearances between wires on distribution lines are less and distribution lines have an array of uninsulated, structure-mounted equipment (Marti 2002). To minimize the risk of raptor electrocutions, TEP will construct the proposed transmission line following the guidelines outlined in “Suggested Practices for Raptor Protection on Power Lines: the State of the Art in 1996” (APLIC 1996). Furthermore, on the structures to be used in the proposed action, the distance between the power lines is at least 18 ft (5.5 m). Because the average wingspan of an adult CFPO is 15 in (38 cm), there is no foreseeable risk of electrocution.

Construction Noise and Activity

Although no CFPO have been detected in the project area, short term noise disturbance and human activity associated with construction may discourage CFPO from using habitat within and adjacent to the proposed ROW. Human activity near nest sites at critical periods of the nesting cycle may cause CFPO to abandon their nests (USFWS 2002a). While CFPO may tolerate low level noise disturbances, such as those in low density residential areas (Cartron et al. 2000b), they will probably not tolerate noise levels associated with construction activities in close proximity to a nest. The greatest likelihood of noise disturbance will result from the use of helicopters during the installation of the transmission lines, but also could result from the presence of heavy machinery or large groups of construction personnel. If CFPO are not detected during the two consecutive years of protocol surveys, the potential for direct impacts to this species is minimal.

Indirect Effects

Habitat Modification and Fragmentation

The proposed action will result in the disturbance of areas that could provide potential nesting, foraging, and dispersal habitat for CFPO. Because many access roads will be closed and restored and all disturbed areas will be reseeded, this disturbance will be temporary. The proposed action could potentially result in temporary disturbance to habitat from access roads and structure installations in the following amounts: 33.99 acres (13.76 ha) in Sonoran desertscrub, 16.70 ha (41.27 acres) in desert riparian scrub, and 0.05 acres (0.02 ha) in deciduous riparian.

While all large saguaros within construction sites will be transplanted, construction could temporarily degrade CFPO habitat by removing vegetation that provides forage and shelter. Elimination of groundcover plant species, rodent burrows, and native soils, as well as loss of trees and shrubs, may impact local reptile and bird populations that are important to the pygmy-owl diet. Loss of complex vegetation structure increases energy demands on owls that must forage at greater distances and risk exposure to a variety of hazards (Abbate et al. 1999). Because of the linear nature of the proposed action, these impacts will be widely distributed and relatively minor in any single area.

Increased Legal and Unauthorized Access to CFPO Habitat

Although CFPO have not been detected in the project area, recreationists may access potential CFPO habitat using temporary construction roads associated with the proposed action. While hikers and other non-motorized recreationists will create minimal disturbance, noise from Off Highway Vehicle (OHV) users are much more likely to disturb CFPO, especially if the activity occurs over an extended period of time in or near a CFPO nesting territory. Increased access to CFPO habitat may subject the species to poaching or other harassment. While TEP will prevent unauthorized access to the ROW across private land, closure of the ROW on public land, particularly state land, is not feasible. Therefore, some increase in access to potential CFPO habitat is anticipated.

Accidental Wildfire

Increased road access may contribute to an increase in the frequency of human caused ignitions in some areas (Gucinski et al. 2001). Because of their mobility, CFPO will not likely be directly impacted by wildfires. However, wildfires may destroy columnar cacti and trees that provide nesting cavities as well as affect CFPO prey species through direct mortality from the fire or habitat destruction. Herbaceous plant species that serve as cover and forage for small mammals could be drastically reduced. Because of reduced groundcover, predation upon surviving small mammals by CFPO may actually increase in the short term. Furthermore, increased herbaceous production in the years following a fire may improve habitat for small mammals in the long term.

New roads also may act as firebreaks and improve response time of firefighters to wildfires, thereby preventing these fires from gaining in size and intensity. A study in southern California concluded that the road network had been a key factor in determining what suppression strategies were used, both in firefighter access and because roads were widely used for backfiring and burning-out operations (Salazar and Gonzalez-Caban 1987). Early studies of fuelbreak effectiveness in southern California came to similar conclusions (Green 1977).

The measures outlined in the Fire Prevention Plan will minimize the risk of wildfire associated with the proposed action.

Invasive Species

Roads may be the first point of entry for invasive species into a new landscape and may serve as a corridor along which plants move farther into the landscape (Lonsdale and Lane 1994, Greenberg et al. 1997). Some invasive plants may then be able to move away from the roadside into adjacent patches of suitable habitat. Invasion by these plants may have significant biological and ecological effects if the species are able to disrupt the structure or function of an ecosystem. Roads constructed for the proposed action could allow the establishment or increased density of non-native plants, such as Lehmann's lovegrass, an invasive species that facilitates wildfires. An increased risk of fire in CFPO habitats could be detrimental to the species because it would eliminate essential features, such as saguaros and desert tree species, which are not fire adapted. Fire stimulates Lehmann's lovegrass, which in turn stimulates more fire, the result is an increase in the fire return interval at the expense of native plant species (McPherson 1995). Measures

outlined in the Invasive Species Management Plan will minimize the introduction or spread of invasive species as a result of the proposed action.

2.1g Cumulative Effects

Cumulative effects include the effects of future state, local, or private actions that are reasonably certain to occur in the action area considered in this biological assessment. While the action area for this species crosses private, state, and federal lands, the habitat with the highest potential for occupancy by CFPO occurs on state and private lands in Pima County. Future federal actions on these lands will be subject to Section 7 consultation. These actions will not be considered cumulative.

Although the amount of future private development within the action area is unknown, many rural areas of Arizona are experiencing substantial growth. Pima County grew by 26.5 percent between 1990 and 2000 (U.S. Census Bureau 2000). Because of the growth rate and the development pressures from nearby Tucson and Sahuarita, it is foreseeable that land adjacent to the proposed ROW will be developed. These developments will likely include increases in associated infrastructure such as roads, groundwater use, and commercial services, all resulting in the degradation of CFPO habitat.

An undetermined level of border crossings by undocumented immigrants (UDI) occurs within the action area, resulting in habitat damage from new roads, discarded trash, illegal campfires, and disturbance near water sources. These border crossings are likely to continue or increase. Additionally, agriculture, recreation, OHV use, grazing, and other activities continue to occur on private and state land and adversely affect CFPO and their habitats.

2.1h Effects Determination and Incidental Take

While CFPO are not currently known to occupy the action area, the disturbance of potential habitat from construction activities and increased access may affect, and are likely to adversely affect, this species.

Take of CFPO is not anticipated because construction activities during breeding season will only occur following protocol surveys and the Conservation Measures outlined in SECTION 1.4 will minimize disturbance to potential habitat and prevent disturbance to nesting CFPO within the action area should any be detected in the future.

2.2 SOUTHWESTERN WILLOW FLYCATCHER (*Empidonax traillii extimus*) (Endangered)

2.2a Action Area

The action area includes all areas potentially affected, directly or indirectly, by all aspects of the project. While habitat for SWFL does not exist within the Central Corridor, migratory SWFL have been documented along the Santa Cruz River, which is within 0.5 mi (0.8 km) of the proposed action near the community of Tumacacori, Arizona.

2.2b Natural History and Distribution

SWFL (Figure 11) are small passerine bird (Order Passeriformes; Family Tyrannidae) measuring approximately 5.75 in (14.6 cm) in length from the tip of the bill to the tip of the tail and weighing only 0.4 ounces (11.34 grams). This species has a grayish-green back and wings, whitish throat, light gray-olive breast, and pale yellowish belly. Two



white wingbars are visible (juveniles have buffy wingbars). The eye ring is faint or absent. The upper mandible is dark and the lower is light yellow grading to black at the tip. SWFL are riparian obligate species, nesting along rivers, streams, and other wetlands where dense growths of willow, seepwillow (*Baccharis* sp.), buttonbush (*Cephalanthus* sp.), boxelder (*Acer negundo*), saltcedar (*Tamarix* spp.), carrizo (*Phragmites australis*) or other plants are present, often with a scattered overstory of cottonwood and/or willow.

Figure 11. Southwestern willow flycatcher.

One of four currently recognized willow flycatcher subspecies (Phillips 1948, Unitt 1987, Browning 1993), SWFL are neotropical migratory species that breed in the southwestern U.S. from approximately 15 May to 1 September. This species migrates to Mexico, Central America, and possibly northern South America during the non-breeding season (Phillips 1948, Stiles and Skutch 1989, Peterson 1990, Ridgely and Tudor 1994, Howell and Webb 1995). The historical range of SWFL included southern California, Arizona, New Mexico, western Texas, southwestern Colorado, southern Utah, extreme southern Nevada, and extreme northwestern Mexico (Sonora and Baja) (Unitt 1987).

SWFL breed in dense riparian habitats from sea level in California to just over 7,000 ft (2,134 m) in Arizona and southwestern Colorado. Historic egg/nest collections and species descriptions throughout SWFL range describe the widespread use of willow for nesting (Phillips 1948, Phillips et al. 1964, Hubbard 1987, Unitt 1987, San Diego Natural History Museum 1995). Currently, SWFL primarily use Geyer willow (*Salix geyeriana*), Goodding willow (*Salix gooddingii*), boxelder, saltcedar, Russian olive (*Elaeagnus angustifolia*), and live oak (*Quercus agrifolia*) for nesting. Other plant species less commonly used for nesting include: buttonbush, black twinberry (*Lonicera involucrata*), cottonwood, white alder (*Alnus rhombifolia*), blackberry (*Rubus ursinus*), carrizo, and stinging nettle (*Urtica* spp.). Nesting SWFL exhibit a strong preference for dense vegetation at the nest site, but high variation and density of vegetation at the patch scale (Hatten et al. 2000). Nesting sites are typically close to the edge of the vegetation patch

and close to water (Allison et al. 2000). Based on the diversity of plant species composition and complexity of habitat structure, four basic nesting habitat types can be described for SWFL: monotypic willow, monotypic exotic, native broadleaf dominated, and mixed native/exotic (Sogge et al. 1997).

Open water, cienegas, marshy seeps, or saturated soil are typically in the vicinity of SWFL territories and nests; SWFL sometimes nest in areas where nesting substrates are in standing water (Maynard 1995, Sferra et al. 1995, 1997). Hydrological conditions at a particular site can vary remarkably in the arid southwest within a season and between years. At some locations, particularly during drier years, water or saturated soil is only present early in the breeding season (i.e., May and part of June). However, the total absence of water or visibly saturated soil has been documented at several sites where the river channel has been modified (e.g. creation of pilot channels), where modification of subsurface flows has occurred (e.g. agricultural runoff), or as a result of changes in river channel configuration after flood events (Spencer et al. 1996). Throughout their range, SWFL arrive on breeding grounds in late April and May (Sogge and Tibbitts 1992, Sogge et al. 1993, Sogge and Tibbitts 1994, Muiznieks et al. 1994, Maynard 1995, Sferra et al. 1995, 1997). Nesting begins in late May and early June, and young fledge from late June typically through mid August, but as late as early September.

SWFL are insectivores, foraging in dense shrub and tree vegetation along rivers, streams, and other wetlands. Flying insects are the most important SWFL prey item; however, they will also glean larvae of non-flying insects from vegetation (Drost et al. 1998). Drost et al. (1998) found that the major prey items of SWFL (in Arizona and Colorado), consisted of true flies (Diptera); ants, bees, and wasps (Hymenoptera), and true bugs (Hemiptera). Other insect prey taxa include leafhoppers (Homoptera: Cicadellidae), dragonflies and damselflies (Odonata); and caterpillars (Lepidoptera larvae). Non-insect prey include spiders (Araneae), sowbugs (Isopoda), and fragments of plant material.

2.2c Critical Habitat

Critical habitat for SWFL was originally designated on 22 July 1997 (USFWS 1997b), but on 11 May 2001, the 10th Circuit Court of Appeals set aside the critical habitat designation and instructed USFWS to issue a new designation in compliance with the court ruling. USFWS is currently soliciting information regarding areas important for the conservation of this species in order to re-propose critical habitat.

2.2d Current Status Statewide

The following status of SWFL in Arizona was summarized from Smith et al. (2002). In 2001, 177 sites covering approximately 139 mi (225 km) of riparian habitat were surveyed for SWFL in Arizona. Sites range from 98 ft (30 m) to 8,802 ft (2,683 m) in elevation and 98.5 ft (30 m) to 10 mi (16.1 km) in length. The mean site length was 1 mi (1.6 km). Fifty-two of the 177 sites were not surveyed according to protocol. This was due to time or funding limitations or because unsuitable SWFL habitat was found during the first survey. Of the 177 sites, 20 had not been previously surveyed. Most new survey sites were located along the Colorado River (n = 9) and Gila River (n = 4). Six hundred

thirty-five resident SWFL were documented within 346 territories at 46 sites. AGFD personnel and statewide cooperators recorded 311 pairs.

SWFL were documented along 11 drainages. The greatest concentrations of SWFL were found at Roosevelt Lake (40 percent) and the Winkelman Study Area (35 percent). Resident SWFL were detected at five sites that had been surveyed at least once in previous years. Resident SWFL were documented in two drainages (Virgin River and Cienega Creek) for the first time since protocol surveys began. No historical occurrence record exists for SWFL along the Virgin River and SWFL have not been reported at Cienega Creek since 1964. These colonizations yield evidence of habitat restoration potential in these drainages that can aid in recovery of the SWFL.

2.2e Environmental Baseline

Deciduous riparian vegetation only occurs within the project area at the Peck Canyon crossing. The canyon and associated riparian area supports ash, walnut, and netleaf hackberry but consists primarily of scattered, individual trees with low understory density (Figure 12). This reach of Peck Canyon is ephemeral and water is probably present only for short periods of time following precipitation events. Semidesert grasslands that are subject to grazing characterize the uplands surrounding Peck Canyon. Because of the patchy habitat and lack of surface water, this area likely will not function as SWFL habitat.



Figure 12. Central Corridor crossing of Peck Canyon.

The Central Corridor also passes within 0.5 mi (0.8 km) of the Santa Cruz River near the community of Tumacacori, Arizona. The riparian vegetation within the Santa Cruz floodplain in this area consists of mature Fremont cottonwood and Goodding willow with a greater than 75 percent canopy cover in most places (Figure 13), as well as a well-developed understory. This reach of the Santa Cruz River is perennial.



Figure 13. Santa Cruz River near Tumacacori, Arizona.

The nearest recent (1999) reports of SWFL are from the Santa Cruz River between Tubac and Rio Rico (McCarthy et al. 1998, Paradzick et al. 1999, Paradzick et al. 2000). All of these reports were of migrant SWFL. Additionally, in May 1998, USFWS personnel located a calling willow flycatcher at the I-19 frontage road across Peck Canyon (USFWS 2001a), 2.5 mi (4 km) east of the proposed action. No follow-up surveys were conducted and it is unknown if this was a migrant or breeding flycatcher.

2.2f Effects of Proposed Action on the SWFL

Direct Effects

Construction Noise and Activity

Noise from helicopter flights associated with construction activities may disturb SWFL using suitable habitat along the Santa Cruz River. However, because I-19 is adjacent to the Santa Cruz River, any SWFL using the river will already be subject to a certain level of ambient noise from traffic. Because of the distance of the proposed action from the

Santa Cruz River and the existing noise level along I-19, any increase in noise associated with the proposed action will be minimal and short term.

Indirect Effects

Habitat Modification and Fragmentation

Deciduous riparian vegetation only occurs within the project area at the Peck Canyon crossing. The proposed action spans Peck Canyon parallel to the existing EPNG gasline and no new access roads are planned within the habitat. This portion of riparian habitat is not suitable for breeding SWFL, therefore, no indirect effects to SWFL through habitat modification are anticipated.

Increased Legal and Unauthorized Access to SWFL Habitat

Access and construction roads for the proposed action will typically be spurs off the existing EPNG gasline and range between 500 – 1,000 ft (152 m and 305 m) in length. Because of the short lengths of the new roads and the presence of I-19 between the proposed action and the Santa Cruz River, there will not be any foreseeable increase in access to SWFL habitat.

Accidental Wildfire

Increased road access may contribute to an increase in the frequency of human-caused ignitions in some areas (Gucinski et al. 2001). New roads also may act as firebreaks and improve response times of firefighters to wildfires, thereby preventing these fires from gaining in size and intensity. A study in southern California concluded that the road network had been a key factor in determining what suppression strategies were used, both in firefighter access and because roads were widely used for backfiring and burning-out operations (Salazar and Gonzalez-Caban 1987). Early studies of fuelbreak effectiveness in southern California came to similar conclusions (Green 1977).

While there is a minimal risk from accidental wildfire associated with the proposed action, any fire will have to spread a significant distance to the east before impacting suitable SWFL habitat. Several roads that could serve as firebreaks and afford firefighting accessibility, most notably I-19, occur between the proposed action and suitable SWFL habitat. Furthermore, measures outlined in the Fire Prevention Plan will minimize the risk of wildfire associated with the proposed action.

Invasive Species

Roads may be the first point of entry for invasive species into a new landscape and may serve as a corridor along which plants move farther into the landscape (Lonsdale and Lane 1994, Greenberg et al. 1997). The short lengths of new access roads, their distance from SWFL habitat, as well as the measures outlined in the Invasive Species Management Plan, will minimize the introduction or spread of invasive species into SWFL habitat.

2.2g Cumulative Effects

Cumulative effects include the effects of future state, local, or private actions that are reasonably certain to occur in the action area considered in this BA. The land between the

proposed action and the Santa Cruz River consists almost exclusively of private land. While Federal actions on these lands will be subject to Section 7 consultation, and therefore not considered cumulative, many private actions could occur without consultation.

Although the amount of planned private development within the action area is unknown, many rural areas of Arizona are experiencing substantial growth. Between 1990 and 2000, Pima County grew by 26.5 percent and Santa Cruz County by 29.3 percent (U.S. Census Bureau 2000). Because of these growth rates and the trend of rural development to occur in areas with some existing infrastructure, it is foreseeable that the private ranches adjacent to Arivaca Road could be sold and subdivided for residential homes and ranchettes. Any substantial population increase in the area also could increase demands for access to recreational land, increase groundwater pumping, and foster the development of commercial services. These impacts to the watershed could degrade the value of habitat within Sopori Wash preventing its use by a variety of species.

An undetermined level of border crossings by UDI occurs within the action area, resulting in habitat damage from new roads, discarded trash, illegal campfires, and disturbance near water sources. These border crossings are likely to continue or increase into the foreseeable future.

2.2h Effects Determination and Incidental Take

Noise from construction of the proposed action may affect SWFL, but it is not likely to adversely affect the species because any increase in noise will be minimal compared to ambient noise levels and short term in duration. Because the proposed action is not likely to adversely affect the species, no take of SWFL is anticipated.

2.3 LESSER LONG-NOSED BAT (*Leptonycteris curasoae yerbabuenae*) (Endangered)

2.3a Action Area

The action area includes all areas potentially affected, directly or indirectly, by all aspects of the project. Potential roosting habitat for LLNB occurs in the Tumacacori and Atascosa/Pajarito mountains, and foraging habitat occurs through those portions of the proposed ROW that contain agave and saguaro cacti. Because LLNB have been documented foraging up to 40 mi (64 km) from roost sites, the action area for the LLNB consists of all potential foraging and roosting habitat within a 40 mi (64 km) buffer surrounding the proposed action.

2.3b Natural History and Distribution

LLNB (formerly Sanborn's long-nosed bat) are one of three members of American leaf-nosed bats (Family Phyllostomidae) in Arizona (Hoffmeister 1986). LLNB (Figure 14) is one of the larger Arizona bats, gray to reddish brown in color. This bat has an erect triangular flap of skin (nose leaf) at the end of a long slender nose. LLNB can be distinguished from *Macrotus* by a much longer nose, greatly reduce tail membrane, and smaller ears; and from *Choeronycteris*, which has a shorter tail, larger tail membrane, and longer, narrower nose than LLNB.



Figure 14. Lesser long-nosed bat.

LLNB occur from the southern United States to northern South America, including several islands and the adjacent mainland of Venezuela and Colombia. LLNB are found between 4 degrees to 32 degrees N latitude in semiarid to arid conditions (Nowak 1994). This bat is typically associated with their primary food source, flower nectar and fruit of columnar cacti, and flower nectar of certain agave species. Because of the seasonal nature of their food source, they must migrate to follow flowering and fruiting plants. In addition to food availability, there must be suitable roosting within commuting distance of the food source. Currently, the longest known commute distance is about 48 km (30 mi).

The primary range of this bat lies in Mexico and Central America. Occurrences in Arizona probably represent range expansion. Prior to the 1930s, there are no records of LLNB in Arizona (Cockrum 1991). Colossal Cave and the Old Mammon Mine are the most northern sites known to house colonies of these bats. However, these sites support colonies of about 5,000 individuals, versus sites in Mexico, which are as large as 150,000 individuals.

LLNB have a bi-seasonal occurrence in Arizona. The maternity season, when bats migrate to southwestern Arizona, represents a United States population of about 30,000 individuals. The other is the fall agave flowering season, located in southeastern Arizona, which attracts about 70,000 bats. Each of these areas contains three known primary roosts and some number of secondary/transient or night roosts (sheltering ten to a few hundred individuals/site).

With the exception of a small bachelor roost located in the Chiricahua Mountains, all remaining records represent small numbers (usually single individuals) at hummingbird feeders, caught in mist nets, or chance findings in residential areas. Constantine (1966) reported two immature females from Maricopa County, one in Phoenix on 30 August 1963 and the other in Glendale on 16 September 1963. The Glendale specimen was found dead. The other was hanging on a screen door (not a normal place) indicating something was likely wrong with that bat. He also reported two males from southern California: one was taken alive on 3 October 1993 outside a home in Yucaipa, the other was taken on 18 October 1996 from the outside of a building in Oceanside (Constantine 1998). LLNB also have been reported from the Aravaipa Canyon area (Cockrum 1991). Hoffmeister (1986) has a record in the Santa Catalina Mountains, but Cockrum (1991) states it was probably a transcription error because the nectar-feeding bats found there belong to the genus *Choeronycteris*. However, Cockrum (1991) does report LLNB from the Santa Catalina Mountains but only once in a mist net set in Sabino Canyon (a female in June).

The diet of LLNB in Arizona consists primarily of the nectar, pollen, and ripe fruit of columnar cacti (particularly saguaro) and agave (e.g., *Agave chrysantha*, *A. deserti*, *A. palmeri*, and *A. parryi*). LLNB have been demonstrated to be a significant pollinator of saguaros, organpipe cacti (*Stenocereus thurberi*), and agaves (Howell and Roth 1981, Alcorn et al. 1962, and McGregor et al. 1962). Generally, LLNB in Arizona forage after dusk to nearly dawn during the months of May through September. In a single night, LLNB will forage well away from their daytime roost sites. In Sonora, Mexico, bats feed on the mainland by night at Bahia Kino and roost by day on Isla Tiburon, 15 to 20 mi (24 to 32 km) away. The closest sizable densities of columnar cacti to LLNB roosts in the Sierra Pinacate, Sonora, Mexico, are found in Organpipe Cactus National Monument in Arizona, about 25 to 30 mi (40 to 48 km) away (Fleming 1991).

In Arizona, females arrive in late March and early April, then migrate northward through Mexico along a “nectar corridor” provided by columnar cacti such as saguaro and organpipe (Fleming 1991). Female LLNB usually arrive in Arizona pregnant and congregate in traditional maternity roosts at lower elevations, feeding primarily on saguaro nectar (Cockrum 1991). Adult males arrive later in the summer and, along with dispersing members of the maternity roosts, usually roost at higher elevations, especially within proximity to significant stands of flowering agave.

LLNB are gregarious and form large maternity colonies that number in the thousands (Hayward and Cockrum 1971, Hoffmeister 1986). All four of the verified LLNB maternity roosts in the United States are found in Arizona (Cockrum 1991). The largest

and most important of the four is found in a mine located in Organpipe Cactus National Monument. About 15,000 LLNB use this mine as a maternity roost. Young are typically born between mid-May and early June (Cockrum 1991, Hayward and Cockrum 1971).

While in the roost during the day, LLNB engage in various activities such as flying, suckling of young, grooming, resting, and interacting with neighbors. LLNB are particularly active during the day and any disturbance, such as aircraft or other human activities, may cause an expenditure of extra energy (Dalton and Dalton 1993, Dalton et al. 1994). Female LLNB gathered in large maternity colonies are particularly vulnerable to disturbances. Maternity colonies are more sensitive because of the vulnerability of nonvolant young, whose recruitment into the population is essential to maintain a viable population.

2.3c Critical Habitat

No critical habitat has been designated for LLNB.

2.3d Current Status Statewide

USFWS listed LLNB as endangered throughout its range in the southwestern United States and Mexico on 30 September 1988 (USFWS 1988). Loss of roost and foraging habitat, as well as direct take of individual bats during animal control programs (particularly in Mexico) have contributed to the current endangered status of the species. All available information on the species through 1994 was summarized in the Lesser Long-nosed Bat Recovery Plan approved in 1997 (Fleming 1994). The Plan indicates that the species is not in danger of extinction in Arizona or Mexico. The species still warrants some protection, as it is vulnerable to human disturbance at roost sites because of its gregarious behavior. There also is particular concern for the protection of forage plants from disturbance or destruction near roost sites.

The primary threats to LLNB populations are agave harvesting and human disturbance of roosting and maternity colonies. Suitable day roosts and suitable concentrations of food plants are the two resources that are crucial to LLNB (Fleming 1995). The USFWS determined that the LLNB was endangered because of the following factors (USFWS 1988):

- A long term decline in population,
- Reports of absence from previously occupied sites
- Decline in the pollination of certain agaves.

Known major roost sites include 16 large roosts in Arizona and Mexico (Fleming 1995). According to surveys conducted in 1992 and 1993, the number of bats estimated to occupy these sites was greater than 200,000. Twelve major maternity roost sites are known from Arizona and Mexico. Disturbance of these roosts, or removal of the food plants associated with them, could lead to the loss of the roosts. Limited numbers of maternity roosts may be the critical factor in the survival of this species.

2.3e Environmental Baseline

LLNB roosts are not known within the proposed corridor, but field surveys did locate small caves and crevices nearby that could serve as LLNB day roosts (HEG 2002, unpublished data). Furthermore, unsurveyed caves, mineshafts, and adits, which may provide suitable roost sites, occur within the Tumacacori-Atascosa mountains. The two closest known LLNB roost sites are the Cave of the Bells in the Santa Rita Mountains, approximately 32 km (20 mi) to the west, and a cave in the Patagonia Mountains, approximately 56 km (35 mi) to the west. Both of these roost sites are within the known flight distance to the proposed action and may utilize the proposed corridor for foraging.

Saguaro cacti occur within the proposed corridor north of Duval Mine Road, and agaves are present in varying densities south of Arivaca Road. While the exact densities of agaves and saguaro cacti were not determined for this BA, CNF estimates that Palmer's agave is widely scattered over 1 million ac (400,000 ha) at densities of 10 to 200 per acre, generally between the elevations of 3,000 ft (914 m) and 6,000 ft (1,829 m) (USFWS 2002b).

The northern portion of the proposed action is primarily undeveloped but does contain some existing electrical distribution lines as well as low density housing developments near Sahuarita Road. The Mission Mine Complex also is located within this section of the project area and the proposed action passes through the Tumacacori EMA of the CNF. Range condition in areas crossed by the proposed action is moderately high with a stable or unknown trend. While agaves have persisted in areas grazed for more than 100 years, mortality through direct herbivory and trampling is known to occur. There is a forest-wide study to determine the effects of livestock grazing on agaves currently underway (USFWS 2001b). Livestock stocking rates for the allotments within the Tumacacori EMA range from 1,320 AUMs in the Peña Blanca Allotment to 2400 AUMs in the Bear Valley Allotment. Allotment Management Plans for Bear Valley and Sardinia Allotments are currently being revised.

2.3f Effects of Proposed Action on LLNB

Direct Effects

Construction Noise and Activity

Although LLNB roosts have not been detected within the proposed corridor, short term noise disturbance and human activity associated with construction activities may disturb LLNB if they are present in undetected roosts adjacent to the proposed corridor. The greatest likelihood of noise disturbance will result from the use of helicopters during the installation, but could also result from the presence of heavy machinery or large groups of construction personnel in close proximity to an undetected roost. The consequences of disturbance to small numbers of LLNB in day roost will be less serious than disturbance of large aggregations of bats at one location.

Indirect Effects

Habitat Modification

Indirect effects to LLNB may result from the potential reduction in forage resources (agaves and saguaro cacti) during construction of temporary access roads or the installation of transmission structures. Because agaves and saguaro cacti are unevenly distributed and the nectar provided by them are seasonally and geographically separated, the loss of significant numbers of either species may alter LLNB foraging patterns and roost selection within the action area. Even if the loss of a high density patch of flowering agaves does not cause the abandonment of a roost, bat survivorship may be reduced through increased foraging flight distances, related energy expenditures, and increased exposure to predators. Because of the linear nature of the proposed action, however, these impacts will be widely distributed and relatively minor in any single area.

Although all agave and saguaro disturbed as a result of the proposed action will be transplanted immediately outside of the construction zone, the long term survival and future flowering of these specimens is uncertain. Agaves are typically easy to cultivate in warm climates with well drained soils (Gentry 1982), but no long term studies of agave transplant survival have been conducted. Transplantation of saguaro is a common practice within southern Arizona, but preliminary results from a 10 year study indicate that smaller saguaros (<16 ft [5 m] tall) are more successfully transplanted than larger saguaros (HEG, unpublished data).

Even in areas where no agave or saguaro presently exist, dormant seeds may be present in the soil. Construction activities associated with the proposed action may compact soil and alter water infiltration, which may prohibit seed germination.

Increased Legal and Unauthorized Access to LLNB Habitat

Because LLNB are sensitive to human disturbance, (to the point of temporarily abandoning a day roost after a single human intrusion) increased human access to roost sites could negatively impact LLNB. The presence of new roads on state land will not likely result in disturbance to undetected roosts because few sites in this area support the rock outcroppings, caves, and mine shafts necessary for LLNB roosts. The greatest potential for undetected roosts occurs on CNF land. The road closures on CNF land outlined in SECTION 1.4 and in the RA (URS 2003) will minimize the probability of increased human access and disturbance of LLNB in undetected roosts in these areas.

Accidental Wildfire

Increased road access may contribute to an increase in the frequency of human caused ignitions in some areas (Gucinski et al. 2001). Agaves in desert grasslands have evolved with fire, but unnaturally high fire frequency and intensity can lead to the decline or elimination of agave populations. Furthermore, agave mortality from fire may affect the abundance and distribution of blooming agaves for a number of years, especially if there is high mortality within certain age and size classes.

New roads also may act as firebreaks and improve the response time of firefighters to wildfires, thereby preventing these fires from gaining in size and intensity. A study in

southern California concluded that the road network had been a key factor in determining what suppression strategies were used, both in firefighter access and because roads were widely used for backfiring and burning-out operations (Salazar and Gonzalez-Caban 1987). Early studies of fuelbreak effectiveness in southern California came to similar conclusions (Green 1977). If deemed appropriate, new roads may allow fuelwood collection in areas currently not accessible, thereby reducing the density of downed, woody material, which is capable of supplying wildfires across the landscape.

The measures outlined in the Fire Prevention Plan will minimize the risks of wildfires associated with the proposed action.

Invasive Species

Roads may be the first point of entry for invasive species into a new landscape and may serve as a corridor along which plants move farther into the landscape (Lonsdale and Lane 1994, Greenberg et al. 1997). Some invasive plants may then be able to move away from the roadside into adjacent patches of suitable habitat. Invasion by these plants may have significant biological and ecological effects if the species are able to disrupt the structure or function of an ecosystem. Roads constructed for the proposed action could allow the establishment or increased density of non-native plants, such as Lehmann's lovegrass, an invasive species that facilitates wildfires. An increased risk of fire in CFPO habitats could be detrimental to the species because it would eliminate essential features, such as saguaros and desert tree species, which are not fire adapted. Fire stimulates Lehmann's lovegrass, which in turn stimulates more fire, the result is an increase in the fire return interval at the expense of native plant species (McPherson 1995). Measures outlined in the Invasive Species Management Plan will minimize the introduction or spread of invasive species as a result of the proposed action.

2.3g Cumulative Effects

Cumulative effects include the effects of future state, local, or private actions that are reasonably certain to occur in the action area considered in this BA. The action area for this species crosses private, state, and federal land. Future federal actions on USFS land will be subject to Section 7 consultation but these actions will not be considered cumulative. Because the action area for this species includes a 40 mi (64 km) buffer, some of the future planned actions on private and state land in southern Pima County and much of Santa Cruz County may be considered cumulative.

Although the amount of this future private development within the action area is unknown, many rural areas of Arizona are experiencing substantial growth. Pima County grew by 26.5 percent between 1990 and 2000 (U.S. Census Bureau 2000). In the same time period, Santa Cruz County grew by 29.3 percent (U.S. Census Bureau 2000).

An undetermined level of border crossings by UDI occurs within the action area resulting in habitat damage from new roads, discarded trash, illegal campfires, and disturbance near water sources. These border crossings are likely to continue or increase. Additionally, agriculture, recreation, OHV use, grazing, and other activities continue to occur on private and state land that adversely affect LLNB and their habitats.

2.3h Incidental Take

The potential disturbance of LLNB in undetected roosts from construction noise and potential mortality of transplanted forage species may affect, and is likely to adversely affect, this species.

No take of LLNB is anticipated as a result of the proposed action for the following reasons. First, noise disturbance will likely impact small numbers of individuals and will be short term in duration, and secondly, changes in agave and saguaro distribution will not be significant in any single location.

2.4 PIMA PINEAPPLE CACTUS (*Coryphantha scheeri* var. *robustispina*) (Endangered)

2.4a Action Area

The action area includes all areas potentially affected, directly or indirectly, by all aspects of the project. Potential habitat for the PPC includes the entire proposed Central Corridor.

2.4b Natural History and Distribution

PPC (Figure 15) are small, round cacti with finger-like projections. Adult cacti range in size from 1.8 in (4.6 cm) to 18 in (46 cm) in height. At the tip of each projection or tubercle is a rosette of 10 to 15 straw-colored spines with one central hooked spine. Plants can be single or multi-stemmed and produce bright yellow flowers after summer rains (Roller 1996).



Figure 15. Pima pineapple cactus.

Populations of PPC are known to occur south of Tucson, in Pima and Santa Cruz counties, Arizona and in adjacent northern Sonora, Mexico. It is distributed at low densities within the Altar and Santa Cruz Valleys, as well as in low lying areas connecting these valleys.

PPC populations are generally found in open patches within semidesert grassland and Sonoran desertscrub plant communities (Brown 1994). They are typically found on flat alluvial bajadas that are comprised of granitic material and are most abundant within the ecotone between the grassland and desertscrub biomes (Roller 1996). This plant is found at elevations between 2,362 (720 m) and 4,593 ft (1,400 m). Typically, PPC are not found in washes or riparian areas.

2.4c Critical Habitat

No critical habitat has been designated for this species.

2.4d Current Status Statewide

USFWS listed PPC as endangered throughout its range on 25 October 1993 (58 FR 49875). Habitat loss and degradation, habitat modification and fragmentation, limited geographic distribution, the rarity of this plant species, illegal collection, and difficulties in protecting areas large enough to maintain functioning populations, all are factors that contribute to the current endangered status of this species. Due to the limited information on PPC population distributions under current habitat conditions, it is difficult to determine the current status of the plant statewide. USFWS has insufficient data to determine if the majority of populations of PPC can be sustained under current reduced and fragmented conditions. PPC densities vary throughout its range with the highest densities occurring south of Tucson through the Santa Cruz Valley (to Amado and surrounding developed parts of Green Valley and Sahuarita, and parts of the San Xavier

District of the Tohono O'odham Nation). Continued urbanization, farm and crop development, mine expansion, and invasion of non-native species are primary threats to PPC populations. Overgrazing by livestock, illegal plant collection, and fire-related interactions involving non-native Lehmann's lovegrass also may have negative impacts on PPC (USFWS 1993).

2.4e Environmental Baseline

The environmental baseline for the PPC evaluates the effects of past and ongoing human and natural factors leading to the current status of the species, its habitat and ecosystem within the action area. Based on monitoring results, the status of the PPC appears to have been recently affected by threats that completely alter or considerably modify more than one-third of the species surveyed habitat and have caused the elimination of nearly 60 percent of documented locations (USFWS 2001c). Dispersed, patchy clusters of individuals are becoming increasingly isolated as urban development, mining, and other commercial activities continue to negatively impact PPC habitat.

The Central Corridor is primarily undeveloped but contains some existing electrical distribution lines and associated roads (Figure 14) and is in close proximity to low density housing developments, and the Mission Mine Complex. A majority of the corridor also parallels the previously disturbed EPNG gasline. While portions of the existing EPNG gasline access road appear relatively unused and support early successional plants (Figure 12), other areas are severely eroded and virtually impassable by motor vehicles.

Surveys for PPC were conducted using an approved survey protocol (Roller 1996) by establishing a belt transect across identified potential habitat with each surveyor covering a 16.4 to 23 ft (5 to 7 m) swath. One survey pass of the entire corridor was conducted with more intensive area searches around confirmed PPC locations. Surveys on state, private, and BLM land covered a 200 ft (61 m) wide area centered on the proposed structure alignment. On the CNF, the coverage was expanded to 750 ft (229 m) wide. All detected PPC locations were recorded using a Global Positioning System (GPS) unit. To determine the extent of proposed disturbance to PPC habitat, recent aerial photography was used to eliminate areas not suitable for PPC, including slopes over 15 percent, high clay or bedrock soils, washes, and previously disturbed areas such as roads, buildings, mining disturbance, etc. During surveys conducted between July 2002 and March 2003, 78 PPC were detected within the 125 ft (38.1 m) ROW between the TEP South Substation and the CNF boundary (HEG 2003, unpublished data). Based on the acreage surveyed, the density of PPC within this area is approximately 0.13 PPC/ac (0.32 PPC/ha).

2.4f Effects of Proposed Action on the PPC

Direct Effects

Because the precise locations of structures and access roads can be modified to avoid sensitive resources, the proposed action will not result in the loss of any individual PPC. All known individual PPC near construction areas and along main access routes will be clearly marked and protected to avoid impacts.

Indirect Effects

Modification of Habitat

The construction of new access roads and the installation of structures will alter PPC seed sources in unoccupied, but potential PPC habitat. Construction vehicles will compact soil, changing water infiltration rates, and road construction will dramatically alter soil structure and seed source depth. Areas around structure sites and many access roads will be temporary and will regenerate as potential PPC habitat in the future. Recent observations indicate that PPC may readily establish in recently disturbed habitats (USFWS 2002c), but these areas must be allowed to recover for years or possibly decades.

Detailed analysis of impacts to habitat for this species is ongoing. To mitigate for the potential loss of PPC habitat, TEP will purchase credits in a USFWS-approved conservation bank for PPC at a ratio determined in consultation with the USFWS.

Increased Legal and Unauthorized Access to PPC Habitat

Much of the proposed corridor through PPC habitat parallels existing electrical distribution lines with existing utility access roads. Some new access roads, however, will be constructed, potentially resulting in unintended access into previously undisturbed PPC habitat (especially by OHV users). Off-road travel could directly impact additional PPC or impede seedling establishment through changes in soil characteristics. Where possible, TEP will review the potential for closure of roads on private land to limit unauthorized access to the ROW.

Accidental Wildfire

Increased road access may contribute to an increase in the frequency of human-caused ignitions in some areas (Gucinski et al. 2001). It is widely regarded that most succulent species are negatively impacted by fire and are not fire adapted (Rogers and Steele 1980, McLaughlin and Bowers 1982). Plants die by direct heating of the fire or later through indirect fire effects such as grazing of spineless plants, post-fire increase in plant tissue temperature, or the introduction of disease or infestation into weakened plants (Thomas 1991). The sparse distribution of this species across the landscape can mean that loss of just a few individuals to fire can greatly affect the range and density of local PPC populations.

New roads may act as natural firebreaks and improve response times of firefighters to wildfires, thereby preventing fires from gaining in size and intensity. A study in southern California concluded that the road network had been a key factor in determining what

suppression strategies were used, both in firefighter access and because roads were widely used for backfiring and burning-out operations (Salazar and Gonzalez-Caban 1987). Early studies of fuelbreak efficacy in southern California came to similar conclusions (Green 1977).

The measures outlined in the Fire Prevention Plan will minimize the risks of wildfires associated with the proposed action.

Invasive Species

Roads may be the first point of entry for invasive species into a new landscape and may serve as a corridor along which plants move farther into the landscape (Lonsdale and Lane 1994, Greenberg et al. 1997). Some invasive plants may then be able to move away from the roadside into adjacent patches of suitable habitat. Invasion by these plants may have significant biological and ecological effects if the species are able to disrupt the structure or function of an ecosystem. Roads constructed for the proposed action could allow the establishment or increased density of non-native plants, such as Lehmann's lovegrass, an invasive species that facilitates wildfires. An increased risk of fire in CFPO habitats could be detrimental to the species because it would eliminate essential features, such as saguaros and desert tree species, which are not fire adapted. Fire stimulates Lehmann's lovegrass, which in turn stimulates more fire, the result is an increase in the fire return interval at the expense of native plant species (McPherson 1995). Measures outlined in the Invasive Species Management Plan will minimize the introduction or spread of invasive species as a result of the proposed action.

2.4g Cumulative Effects

Cumulative effects include the effects of future state, local, or private actions that are reasonably certain to occur in the action area considered in this biological assessment. Under Section 9 of the Act, the taking of listed animals is specifically prohibited, regardless of land ownership status. For listed plants, these prohibitions and the protection they afford do not apply. Listed plant species are protected only from deliberate removal from Federal land. There is no protection against removal or destruction of plants by a landowner on private land under the ESA.

Although the amount of future private development within the action area is unknown, many rural areas of Arizona are experiencing substantial growth. Pima County grew by 26.5 percent between 1990 and 2000 (U.S. Census Bureau 2000). Because of these growth rates and the development pressures of nearby Tucson and Sahuarita, Arizona, it is foreseeable that some lands adjacent to the proposed ROW will be developed. These developments will likely include increases in associated infrastructure such as roads, groundwater use, and commercial services, all resulting in the degradation of PPC habitat.

An undetermined level of border crossings by UDI occurs within the action area and results in habitat damage from new roads, discarded trash, illegal campfires, and disturbance near water sources. These border crossings are likely to continue or increase.

Additionally, PPC habitat is adversely affected by continual agriculture, recreation, OHV use, grazing, and other activities on private and state land.

2.4h Effects Determination

Construction activities and increased access may affect, and are likely to adversely affect PPC within the ROW, potential PPC habitat, and seedling establishment. The adverse affects to the species will be mitigated through the purchase of mitigation bank credits.

2.5 JAGUAR (*Panthera onca*) (Endangered)

2.5a Action Area

The action area includes all areas potentially affected, directly or indirectly, by all aspects of the project. Because of the large movements possible by the jaguar and historical records for the species in a variety of habitats, the action area for the jaguar considered for the proposed action includes most of western Santa Cruz and southern Pima counties.

2.5b Natural History and Distribution

Jaguars (Figure 16) are the largest species of cat now native to the Western Hemisphere. Jaguars are large muscular cats with relatively short massive limbs, a deep-chested body, and cinnamon-buff in color with many black spots. Its range in North America includes Mexico and portions of the southwestern United States (Hall 1981). A number of jaguar records are known for Arizona, New Mexico, and Texas. Additional reports exist for California and Louisiana. Records of the jaguar in Arizona and New Mexico have been attributed to the subspecies *Panthera onca arizonensis*. The type specimen of this subspecies was collected in Navajo County, Arizona, in 1924 (Goldman 1932). Nelson and Goldman (1933) described the distribution of this subspecies as the mountainous parts of eastern Arizona north to the Grand Canyon, the southern half of western New Mexico, northeastern Sonora, and, formerly, southeastern California. The records for Texas have been attributed to another subspecies *P. o. veraecrucis*. Distribution of this subspecies was described by Nelson and Goldman (1933) as the Gulf slope of eastern and southeastern Mexico from the coast region of Tabasco, north through Vera Cruz and Tamaulipas, to central Texas. Swank and Teer (1989) indicated the historical range of the jaguar included portions of Arizona, New Mexico, and Texas. These authors consider the current range to be central Mexico through Central America and into South America as far as northern Argentina.




Figure 16. Jaguar.

Swank and Teer (1989) stated the United States no longer contains established breeding populations of jaguar, which probably disappeared from the United States in the 1960s. According to these authors, the jaguar prefers a warm tropical climate and is usually associated with water, and rarely found in extensive arid areas. Goldman (1932) believed the jaguar was a regular, but not abundant, resident in southeastern Arizona. Hoffmeister (1986) considered the jaguar an uncommon resident species in Arizona. He concluded that the reports of jaguars between 1885 and 1965 indicated a small but resident population once occurred in southeastern Arizona. Brown (1983a) suggested the jaguar in Arizona ranged widely throughout a variety of habitats from Sonoran desert scrub through subalpine conifer forest. Most of the records were from Madrean evergreen-woodland, shrub-invaded semidesert grassland, and along rivers.

Brown (1983a) presented an analysis suggesting there was a resident breeding population of jaguars in the southwestern United States at least into the 20th century. USFWS (1990) recognized that the jaguar continues to occur in the American southwest as an occasional wanderer from Mexico. Currently, breeding population of jaguar are unknown in the United States.

In Arizona, the gradual decline of the jaguar appeared to be concurrent with predator control associated with land settlement and the development of the cattle industry (Brown 1983a, USFWS 1990). Lange (1960) summarized the jaguar records from Arizona, and between 1885 and 1959 the reports consisted of 45 jaguars killed, six sighted, and two recorded by sign. Brown (1991) related that the accumulation of all known records indicated a minimum of 64 jaguars were killed in Arizona after 1900.

2.5c Critical Habitat

No critical habitat has been designated for this species.

2.5d Current Status Statewide

Jaguar were initially listed as endangered from the United States - Mexico border southward to include Mexico and Central and South America (37 FR 6476, 1972; 50 CFR 17.11, August 1994). As a result of a petition, the jaguar was proposed as endangered in the United States (59 FR 35674; July 13, 1994). In a Federal Register notice dated 22 July 1997, the jaguar was listed as an endangered species in the United States (62 FR 39147).

The most recent records of jaguars in the United States are from Arizona. In 1971, a jaguar was taken east of Nogales and in 1986 one was taken from the Dos Cabezas Mountains. The latter reportedly had been in the area for about a year before it was killed. AGFD (1988) cited two recent reports of jaguars in Arizona. The individuals were considered to be transients from Mexico. One report (1987) was from an undisclosed location. The other report was from 1988, when tracks were observed for several days prior to the treeing of a jaguar by hounds in the Altar Valley, Pima County. An unconfirmed report of a jaguar at the Coronado National Memorial was made in 1991. In 1993, an unconfirmed sighting of a jaguar was reported for Buenos Aires National Wildlife Refuge. In March 1996, the presence of a jaguar was confirmed through photographs made in the Peloncillo Mountains of Arizona and New Mexico (Glenn 1996). AGFD reported a jaguar sighting in the Baboquíviri Mountains in 1996, and in the fall of 1997, one was reported from the Cerro Colorado Mountains of southern Arizona. A jaguar was recently documented (December 2001) in the Atascosa Mountains within about 2 mi (3 km) of the proposed action.

2.5e Environmental Baseline

The Tumacacori EMA is the location of recent reports of jaguars in the United States. This area continues to include the most likely habitat that will support the existence of jaguars in the United States. Many of the larger canyon bottoms in the Tumacacori EMA contain substantial cover and could act as travel corridors for dispersing jaguars. It is believed that all recent sightings of jaguars in Arizona are males dispersing north from

the northern most breeding population in Mexico in an effort to find unoccupied habitat (B. VanPelt, AGFD, pers. comm., 3 October 2002). Because no breeding pairs are thought to exist north of the United States-Mexico border, conservation of the Mexican population is vital to the future presence of jaguars in Arizona.

Under the leadership of AGFD and New Mexico Department of Game and Fish, a conservation agreement and strategy has been prepared to address the conservation of the jaguar in Arizona and New Mexico. This agreement established an interstate/intergovernmental Jaguar Conservation Team under a Memorandum of Agreement (MOA). This MOA has been signed by various state and federal cooperators and local and tribal governments with land and wildlife management responsibilities in the geographic area of concern. The Jaguar Conservation Agreement and Strategy serves as a mechanism for implementation of actions for the protection and conservation of the jaguar, while providing a template for the recovery of the species until a recovery plan is prepared and adopted.

The Conservation Agreement established procedures for reporting and evaluating jaguar sightings and compiling distribution and occurrence information, investigation of livestock depredation, evaluation of habitat suitability, development of education materials, and other activities. The Jaguar Conservation Agreement also provides for participation by interested private citizens and organizations. CNF grazing allotment permittees are participating in this process.

The December 2001 sighting mentioned earlier came from a remote camera operated under the direction of the Jaguar Conservation Team (S. Schwartz, AGFD, pers. comm., 17 September 2002). Currently, 14 remote cameras are positioned along the United States-Mexico border in an attempt to document movement of jaguars in and out of Arizona (J. Childs, Jaguar Conservation Team, pers. comm., 3 October 2002).

2.5f Effects of Proposed Action on the Jaguar

Direct Effects

Construction Noise and Activity

Because jaguars are primarily nocturnal, disturbance from construction activities, even in suitable dispersal habitat, is unlikely. The greatest likelihood of noise disturbance will result from the use of helicopters during early morning or late evening hours. However, because of the linear nature of the proposed action, any noise disturbance will be widely distributed and relatively short term in any location. Any jaguar within the action area will likely avoid construction sites. The use of additional remote cameras to monitor the United States-Mexico border south of the proposed action also will minimize the possibility of construction activities affecting breeding jaguars.

Indirect Effects

Habitat Modification and Fragmentation

Roads can reduce habitat value because of habitat fragmentation and edge effects. Some studies have shown that a few large areas of low road density, even in a landscape of high

average road density, may be the best indicator of suitable habitat for large vertebrates (Rudis 1995). Because construction activities within riparian corridors or other major canyons will be minimal and widely distributed, no adverse impacts to the composition or structure of jaguar movement corridors or fragmentation of habitat is anticipated. Furthermore, access and construction roads for the proposed action commonly are spurs off existing roads and range between 500 ft (152 m) and 1,000 ft (305 m) in length, which do not isolate or separate habitat patches.

While access roads and structure site construction could degrade the habitats of jaguar prey species, effects on the prey base are difficult to quantify. The primary jaguar prey species in Arizona is deer (*Odocoileus* spp.), which have relatively large home ranges. Road-avoidance behavior (up to distances of 300 ft [90 m] to 600 ft [180 m]) is common in large mammals (Lyon 1983), including those species that may serve as prey for jaguars. Because of the linear nature of the proposed action, impacts to deer habitat will be widely distributed and relatively minor in any single area.

Increased Legal and Unauthorized Access to Jaguar Habitat

Jaguars appear to be relatively tolerant of some level of human activity (B. VanPelt, AGFD, pers. comm., 3 October 2002) and have been documented using areas that have recreational and agricultural activities occurring on a regular basis. However, increased human access to potential jaguar habitat through the use of temporary proposed construction roads could reduce the quality of the habitat. The road closure techniques outlined in the SECTION 1.4 and the RA (URS 2003) will minimize unintended uses of these roads.

Accidental Wildfire

Increased road access may contribute to an increase in the frequency of human-caused ignitions in some areas (Gucinski et al. 2001). Because of their mobility, jaguars will not likely be directly impacted by wildfires; however, these wildfires could potentially alter or destroy portions of prey species habitat. While the short-term effects of wildfires may affect prey species through loss of forage from the fire, increased herbaceous production in the years following a fire may improve habitat in the long term.

New roads also may act as firebreaks and improve response time of firefighters to wildfires, thereby preventing these fires from gaining in size and intensity. A study in southern California concluded that the road network had been a key factor in determining what suppression strategies were used, both in firefighter access and because roads were widely used for backfiring and burning-out operations (Salazar and Gonzalez-Caban 1987). Early studies of fuelbreak effectiveness in southern California came to similar conclusions (Green 1977). If deemed appropriate, new roads may allow fuelwood collection in areas currently not accessible, thereby reducing the density of downed, woody material, which is capable of carrying wildfires across the landscape. The fire prevention measures being developed for the Fire Prevention Plan will minimize the risks of wildfires associated with the proposed action.

Invasive Species

Roads may be the first point of entry for invasive species into a new landscape, and can serve as a corridor along which plants move farther into the landscape (Lonsdale and Lane 1994, Greenberg et al. 1997). Some invasive plants may then be able to move into adjacent patches of suitable habitat. Invasion by these plants may have significant biological and ecological effects if the species are able to disrupt the structure or function of an ecosystem. Roads constructed for the proposed action could allow the establishment or increased density of non-native plants, such as Lehmann's lovegrass, an invasive species that facilitates wildfires (McPherson 1995). Measures outlined in the Invasive Species Management Plan will minimize the introduction or spread of invasive species as a result of the proposed action.

2.5g Cumulative Effects

Cumulative effects include the effects of future state, local, or private actions that are reasonably certain to occur in the action area considered in this BA. While the action area for this species encompasses private, state, and federal lands, the habitat with the highest potential for occupancy by jaguars occurs on USFS land in Santa Cruz County. Future federal actions on these lands will be subject to Section 7 consultation; these actions will not be considered cumulative.

Although the amount of future private development within Santa Cruz County is unknown, many rural areas of Arizona are experiencing substantial growth. Between 1990 and 2000, Santa Cruz County grew by 29.3 percent (U.S. Census Bureau 2000). Despite its distance from the proposed action, an increase in population in Nogales, Arizona and other regional population centers translates into an increased demand recreational use of USFS land.

An undetermined level of border crossings by UDI also occurs within the action area, resulting in habitat damage from new roads, discarded trash, illegal campfires, and disturbance near water sources. These border crossings are likely to continue or increase.

2.5h Effects Determination and Incidental Take

Construction noise and activity associated with the proposed action may affect the jaguar, but it is not likely to adversely affect the species because any disturbance will be widely distributed and short term in duration.

Because the proposed action is not likely to adversely affect the jaguar, no take is anticipated.

2.6 GILA TOPMINNOW (*Poeciliopsis occidentalis occidentalis*) (Endangered)

2.6a Action Area

The action area includes all areas to be affected directly or indirectly by the federal action and not merely the immediate area involved in the action. In streams, the action area is often much larger than the area of the proposed action because impacts in the watershed may be concentrated in the stream and actions within the stream may be carried downstream well outside of the immediate project area. The action area for the Gila topminnow is the entire Santa Cruz River watershed.

2.6b Natural History and Distribution

The Gila topminnow (Figure 17) was originally described by Baird and Girard (1853) as *Heterandria occidentalis* from a specimen collected in 1851 from the Santa Cruz River near Tucson. It was redescribed by Hubbs and Miller (1941) as *Poeciliopsis occidentalis*. As with all species in the family Poeciliidae, the Gila topminnow exhibits sexual dimorphism. Both males and females are tan to olive-bodied and usually white on the belly. Scales of the dorsum are darkly outlined and the fin rays contain melanophores, although lacking in dark spots. Dominant sexually mature males are often blackened,



Figure 17. Gila topminnow

with some gold on the pre-dorsal midline, orange at the base of the gonopodium, and exhibits bright yellow pelvic, pectoral, and caudal fins (Minckley 1973). Females remain drab in coloration upon reaching maturity and throughout their life. All male poeciliids have a modified anal fin (gonopodium) used to fertilize the female internally.

Habitat requirements of *P. o. occidentalis* are broad. The species prefers shallow, warm, fairly quiet water; however, they can become acclimated to a much wider range of conditions. Both lentic habitats and lotic habitats with moderate current are easily tolerated. Temperatures from near freezing under ice to 98.6 degrees F (37 degrees C) have been reported, with a maximum tolerance of 109.4 degrees F (43 degrees C) for brief periods (Heath 1962). Gila topminnows can live in a wide range of water chemistries, with recorded pH values from 6.6 to 8.9, dissolved oxygen readings from 2.2 to 11 milligrams/liter (Meffe et al. 1983), and salinities from very dilute to sea water (Schoenherr 1974). The widespread historic distribution of Gila topminnows throughout rivers, streams, marshes, and springs of the Gila River Basin is evidence for their tolerance of these environmental extremes. One reestablished population (Mud Springs) survived for 16 years in a simple cement-watering trough before being moved.

Meffe et al. (1983) reported that topminnows can tolerate almost total loss of water by burrowing into the mud for 1-2 days. Preferred habitats contain dense mats of algae and debris, usually along stream margins or below riffles, with sandy substrates sometimes covered with organic mud and debris (Minckley 1973). Topminnows are usually found in the upper third of the water column and young show a preference for the warmest and shallowest areas (Forrest 1992). Simms and Simms (1992) found topminnows occupying pools, glides, and backwaters more frequently than marshes or areas of fast flow.

According to Schoenherr (1974), the spring-heads presently occupied by Gila topminnows are questionable as preferred habitat. Destruction of historically occupied habitats such as the marshes, sloughs, backwaters, and edgewaters of larger rivers and presence of non-native fish in such habitats that remain has undoubtedly forced Gila topminnow out of their preferred historic habitats and into the spring-heads and smaller erosive creeks we see them in today. Their tolerance of conditions in these habitats has allowed them to maintain populations with less impact from non-native fishes.

Gila topminnows are viviparous fish, meaning embryos grow and mature within the female and are born living. Eggs are fertilized internally through deposition of spermatophores (packets of sperm) into the female genital pore by the male gonopodium. Female Gila topminnow can store spermatozoa for several months, and may produce up to 10 broods after being isolated from males (Schultz 1961). Female Gila topminnows also exhibit superfetation in which 2 or more groups of embryos at different stages develop simultaneously. Females of the genus *Poeciliopsis* generally carry only 2 stages, although some *P. o. occidentalis* females have been shown to carry 3 stages for a few days when population densities are low. The mean interval between broods is 21.5 days (Schoenherr 1974). Brood size ranges from 1-31 dependent upon female standard length (SL) (Constantz 1974; Schoenherr 1974, 1977). Under optimum laboratory conditions, *Poeciliopsis* can produce 10 broods per year at intervals of 7 to 14 days (Schultz 1961). Sexual maturity can be attained as early as 2 months or as late as 11 months following birth, dependent upon the season of birth (Schultz 1961; Constantz 1976, 1979; Schoenherr 1974).

Breeding occurs primarily during January through August, but in thermally constant springs, young may be produced throughout the year (Heath 1962; Minckley 1973; Schoenherr 1974). During the peak of the breeding season up to 98 percent of mature females are pregnant (Minckley 1973). Dominant males turn black, defend territories, and court females. Smaller subordinate males do not turn black or defend territories. Instead, they take on a "sneaking" mating strategy where they attempt to mate with uncooperative females while the dominant male is busy elsewhere. Subordinate males have a longer gonopodium, which may have an adaptive benefit for this type of mating strategy (Constantz 1989). However, if the larger territorial males are removed, smaller males will become dominant, take on breeding coloration, and defend territories (Constantz 1975; Schoenherr 1977). Brood size and the onset of breeding in topminnows can be influenced by several factors including food abundance, photoperiod, temperature, predation upon the population, and female size. Increased food supply and larger female size are believed to contribute to the greater fecundity seen in topminnows from Monkey Spring canal compared with topminnows from Monkey Spring headspring (Constantz 1974, 1979; Schoenherr 1974, 1977). Sex ratios in stabilized populations nearly always favor females, varying from 1.5 to 6.3 per male (Schoenherr 1974).

Gila topminnows are opportunistic omnivorous feeders, having a gut length 1.5 to 2 times SL of the individual (Schoenherr 1974). They have weakly spatulate dentition characteristic of an omnivorous diet. Primary food items include detritus, vegetation,

amphipods, ostracods, insect larvae, and rarely, other fish (Schoenherr 1974; Gerking and Plantz 1980; Meffe et al. 1983; Meffe 1984).

Gerking and Plantz (1980) noted that Gila topminnows prefer to eat large prey, but prey sizes are limited by mouth size. Schoenherr (1974) observed that individual fishes in complex habitats with several food resources present will select and focus on different items. He suggested that variation in feeding among individuals prevents over-utilization of a single resource, thus enhancing survival potential of the species.

In the United States, this species currently occurs in the Gila River drainage, Arizona, particularly in the upper Santa Cruz River, Sonoita and Cienega creeks, and the middle Gila River. The Gila topminnow is restricted to 14 natural localities in Arizona. In Mexico, the species occurs in the Río Sonora, Río de la Concepción, and Santa Cruz River but are not listed under the ESA. Gila topminnows occupy a variety of habitats, including: springs, cienegas, permanent and interrupted streams, and margins of large rivers. Habitat alteration and destruction, and introduction of predatory non-native fish, (principally western mosquitofish [*Gambusia affini*]) is the main reason for decline of the Gila topminnow.

2.6c Critical Habitat

No critical habitat has been designated for this species.

2.6d Current Status Statewide

The United States population of the Gila topminnow was federally listed as an endangered species in 1967 (USDOI 1967). The original recovery plan for Gila topminnow listed 10 extant natural populations: Monkey Spring, Cottonwood Spring, Sheehy Spring, Sharp Spring, Santa Cruz River near Lochiel, Redrock Canyon, Cienega Creek, Sonoita Creek (presumably including localities above and below Patagonia Lake), Salt Creek, and Bylas Springs (USFWS 1984). Gila topminnows were also known from Middle Spring (also known as SII or Second Spring) on the San Carlos Apache Indian Reservation (Meffe et al. 1983). Middle Spring was considered part of the Bylas Springs complex in the earlier recovery plan.

Since 1984, Gila topminnows have been discovered or rediscovered at 4 additional locations: North Fork of Ash Creek in 1985 (Jennings 1987), Fresno Canyon in 1992, Santa Cruz River north of Nogales in 1994, and Coal Mine Canyon in 1996 (Weedman and Young 1997). However, Gila topminnow were last collected from the North Fork of Ash Creek in 1985 and from Sheehy Spring in 1987. They have also been very rare or absent during recent surveys (last 5 years) of Sonoita Creek above Patagonia Lake and Santa Cruz River near Lochiel. Mosquitofish are quite common in both areas. Topminnows were extirpated from 1 of the original 10 localities, Salt Creek, by mosquitofish (Marsh and Minckley 1990), but the stream was renovated and restocked with Gila topminnows from Middle Spring. Subsequently, mosquitofish were found in the stream and it was again renovated and restocked with topminnows from Bylas Spring. Thus, there are 14 naturally occurring localities (considering Sonoita Creek above and

below Patagonia Lake as 2 separate localities) currently known to support Gila topminnows in the United States.

Eleven of the naturally occurring locations currently supporting Gila topminnows are in the Santa Cruz River system: Redrock Canyon, Cottonwood Spring, Monkey Spring, upper Sonoita Creek, Fresno Canyon, Coal Mine Canyon, lower Sonoita Creek, Santa Cruz River north of Nogales, Cienega Creek, Sharp Spring, and the upper Santa Cruz River. The 2 remaining localities (Bylas Springs and Middle Spring) and Salt Creek are next to the Gila River on the San Carlos Apache Indian Reservation. Bylas Springs has been unsuccessfully poisoned twice to remove mosquitofish (Meffe et al. 1983; Brooks 1985; Marsh and Minckley 1990). Another attempt at renovation of Bylas Springs was done by USFWS Arizona Fishery Resource Office and has so far been successful. The population at Middle Spring was eliminated by lack of water during the summer of 1989, but was recently reestablished (following construction of additional pool habitat) with Gila topminnows from the original Middle Spring population held at Roper Lake State Park. Salt Creek has also been renovated and restocked with topminnows originally from Bylas Spring.

As part of past recovery actions, more than 200 Gila topminnow reintroductions or natural dispersals from reintroductions have occurred at 175 wild locations. For this count, a wild location refers to an area that does not have a mailing address, in contrast with a captive population that does (following Simons 1987). Eighteen wild populations remained in 1997, 17 of which are in historic range (Weedman and Young 1997). Seven of these populations are secure enough that they should persist into the foreseeable future. Minckley and Brooks (1985), Brooks (1985, 1986), Simons (1987), Bagley et al. (1991), Brown and Abarca (1992), and Weedman and Young (1997) describe the plight of re-established and captive populations of Gila topminnows.

Gila topminnows also have been stocked into many captive locations for propagation or conservation. Twelve captive populations were known to persist in 1997. The following publicly maintained populations are large enough to provide individuals for reintroductions, although one is known to be mixed with topminnows from more than one natural population (Arizona-Sonora Desert Museum, Boyce-Thompson Arboretum (mixed), Dexter National Fish Hatchery and Technology Center, Roper Lake State Park, Arizona State University, and Hassayampa River Preserve).

2.6e Environmental Baseline

Gila topminnow currently occupy the Santa Cruz River in its perennial reaches, as far north as Chavez Siding Road. This reach of the river was also occupied by longfin dace (*Agosia chrysogaster*), desert sucker (*Catostomus clarki*), Sonora sucker (*Catostomus insignis*), green sunfish (*Lepomis cyanellus*), and mosquitofish as recently as 1997 (USFWS 2001d). No Gila topminnows occur on the Tumacacori EMA and there are currently no plans for reintroductions in any locations (CNF 2000; D. Duncan, USFWS, pers. comm., 1 October 2002).

2.6f Effects of Proposed Action on the Gila topminnow

Direct Effects

The effects of the proposed action on this species are not anticipated to include direct effects to individual Gila topminnow because no construction will occur within occupied habitat.

Indirect Effects

Habitat Modification

Some indirect impacts to Gila topminnow habitat from erosion are possible from the construction of the proposed action. While the removal of vegetation for construction of access roads will increase surface runoff and sediment transport, and decrease infiltration of precipitation (Gifford and Hawkins 1978, Busby and Gifford 1981, Blackburn 1984, DeBano and Schmidt 1989, Belnap 1992, Belsky and Blumenthal 1997), the implementation of BMPs will help control erosion. However, unusually large precipitation events may temporarily overwhelm BMPs and result in some increase in sediment transport. Nevertheless, the distance of the proposed action from the Santa Cruz River will minimize the amount of sediments reaching Gila topminnow habitat.

Accidental Wildfire

Increased road access may contribute to an increase in the frequency of human-caused ignitions in some areas (Gucinski et al. 2001). Roads constructed for the proposed action also may allow the establishment or increased density of non-native grasses, such as Lehmann's lovegrass, an invasive species that facilitates wildfires (McPherson 1995). Wildfires could remove groundcover that is important in dissipating rainfall energy and reducing erosion.

However, new roads also may act as firebreaks and improve response time of firefighters to wildfires, thereby preventing these fires from gaining in size and intensity. A study in southern California concluded that the road network had been a key factor in determining what suppression strategies were used, both in firefighter access and because roads were widely used for backfiring and burning-out operations (Salazar and Gonzalez-Caban 1987). Early studies of fuelbreak effectiveness in southern California came to similar conclusions (Green 1977). If deemed appropriate, new roads may allow fuelwood collection in areas currently not accessible, thereby reducing the density of downed, woody material, which is capable of carrying wildfires across the landscape.

The measures outlined in the Fire Prevention Plan being developed will minimize the risks of wildfires associated with the proposed action. Measures outlined in the Invasive Species Management Plan also will minimize the introduction or spread of invasive species that may facilitate fires.

2.6g Cumulative Effects

Cumulative effects include the effects of future state, local, or private actions that are reasonably certain to occur in the action area considered in this BA. While the action area for this species encompasses private, state, and federal land, the habitat with the

highest potential for occupancy by Gila topminnow occurs on private land in Santa Cruz County. Most future actions on private land will not be subject to Section 7 consultation.

Although the amount of future private development within Santa Cruz County is unknown, many rural areas of Arizona are experiencing substantial growth. Between 1990 and 2000, Santa Cruz County grew by 29.3 percent (U.S. Census Bureau 2000). Despite its distance from the proposed action, an increase in population in Nogales, Arizona and other regional population centers translates into an increased demand for recreational use of national forest lands.

An undetermined level of border crossings by UDI also occurs within the action area, resulting in habitat damage from new roads, discarded trash, illegal campfires, and disturbance near water sources. These border crossings are likely to continue or increase.

2.6h Effects Determination and Incidental Take

The transport of sediments into the Santa Cruz River may affect the Gila topminnow; however, any increase in sediments will be relatively small because of the distance of the proposed action from occupied habitat. Therefore, it is not likely to adversely affect the species.

Because the proposed action is not likely to adversely affect the species, no take of Gila topminnow is anticipated.

2.7 MEXICAN GRAY WOLF (*Canis lupus baileyi*) (Endangered)

2.7a. Action Area

The action area includes all areas potentially affected, directly or indirectly, by all aspects of the project. Potential habitat for Mexican gray wolf is found within portions of Santa Cruz County containing oak and pine-juniper savannas above 4,000 ft (1,200 m). Wolves may travel long distances during hunting expeditions, typically in an irregular circle 20 mi (34 km) to 60 mi (68 km) in diameter. The action area for the Mexican gray wolf considered for the proposed action includes all potential habitat and travel corridors in western Santa Cruz and southern Pima County.

2.7b. Natural History and Distribution

Mexican gray wolves (Figure 18) are the smallest and southernmost of the 5 subspecies of gray wolf in North America. The Mexican gray wolf is a large dog-like carnivore with a mixed brown, rust, black, gray, and white. This species has a distinct white lip line, chin, and throat. Adults weigh between 50-90 lbs (23-41 kg) (Hoffmeister 1986). The historic range was from southeastern Arizona, southwestern New Mexico, southwestern



Texas, and south through the Sierra Madre of Mexico. The Mexican gray wolf is the southernmost occurring and most endangered subspecies in North America. This wolf is the last subspecies of gray wolf known to occur in the Arizona-New Mexico area. The last known naturally occurring specimen in the United States was found in New Mexico in 1970 (USFWS 2001d).

Figure 18. Mexican gray wolf.

Historically, Mexican gray wolf habitat was montane woodlands, presumably because of the favorable combination of cover, water, and prey availability. Most wolf collections came from pine, oak, and pinyon-juniper woodlands, and intervening or adjacent grasslands above 1,372 m (4,500 ft) (Brown 1983b). Wolves avoided desertscrub and semidesert grasslands, but wooded riparian corridors were probably used for travelling and hunting (Parsons 1996).

These are social animals in the dog family that live and travel in packs of 7 to 30 animals depending upon prey size and availability. Mexican gray wolves prey upon a variety of animals from mice and squirrels to deer and elk. Territory size can range from 30 (78 km²) to 500 mi² (1,295 km²) or more. Packs are led by a pair of dominant animals that control most of the breeding. Breeding season lasts from late winter to early spring, and the dominant female produces up to 6 pups for the pack. The wolves care for the pups communally.

During the late 1800s through the mid 1900s, extensive hunting, trapping, and poisoning efforts at local, state, and federal levels resulted in the extirpation of this species from the United States portion of its range. Reintroduction efforts of captive-bred wolves are under way in the Blue Range Recovery Area of eastern Arizona and New Mexico. Fourteen packs have been released to date.

2.7c Critical Habitat

No critical habitat has been designated for this species.

2.7d Current Status Statewide

Mexican gray wolves were listed as endangered by USFWS in 1976 (41 FR 17736) without critical habitat. In 1998, an experimental, non-essential population was designated for the southwest (63 FR 1763) and a reintroduction program was initiated. Eleven wolves from captive breed stock were reintroduced into the Apache National Forest in southeastern Arizona under the experimental, non-essential designation in an effort to re-establish the subspecies to a portion of its historic range. A Recovery Plan for this subspecies was completed in 1982 and revisions are currently in progress (USFWS 2001d).

Mexican gray wolf populations steadily declined in Arizona because of predator control programs and conflicts with livestock interests. Pressure to control wolves became a priority beginning in the 1920s when this subspecies was nearly eliminated from the state and prevention of wolves from entering from Mexico was undertaken. In 1921 and 1922, a reported 58 wolves were taken by trapping or poisoning in Arizona. By 1924, reported takings dropped to 29 and by 1936, to 5. After 1952, only 2 wolves were reported taken in Arizona, 1 in 1958 and another in 1960 (Hoffmeister 1986). Reports of Mexican gray wolves living in the wild in Arizona continued into the early 1970s (USFWS 1982).

Similar predator control programs in Mexico reduced populations and may have eliminated the wolf by the 1980s. Surveys conducted in Mexico in the early 1990s did not confirm Mexican gray wolf populations in the wild (Parsons 1996).

2.7e Environmental Baseline

The environmental baseline is an analysis of the effects of past and ongoing human and natural factors leading to the current status of the species, its habitat, and ecosystem within the action area. The environmental baseline defines the current status of the species and its habitat in the action area to provide a platform to assess the effects of the action now under consideration.

The Tumacacori EMA contains some areas of montane and riparian woodlands that may serve as dispersal corridors for Mexican gray wolves. If wolf populations exist in the mountains of Sonora, these corridors may be used as hunting and dispersal corridors. There are currently no plans to reintroduce the Mexican gray wolf into southern Arizona and, because of the distance and fragmentation of intervening habitat, it is unlikely that current experimental populations in northern Arizona could disperse into Santa Cruz County.

2.7f Effects of Proposed Action on the Mexican Gray Wolf

Direct Effects

Construction Noise and Activity

Because the only wild populations of Mexican gray wolves in Arizona occur in the Apache National Forest, disturbance from construction of the proposed action, even in suitable dispersal habitat, is highly unlikely. In the event that populations of wolves exist in Mexico and could disperse into southern Arizona, the greatest likelihood of disturbance will result from the use of helicopters during early morning or late evening hours. However, because of the linear nature of the proposed action, any noise or construction disturbance will be widely distributed and relatively minor in any single area.

Indirect Effects

Habitat Modification and Fragmentation

Roads can reduce habitat value because of habitat fragmentation and edge effects. Gray wolves (*Canis lupus*) in Wisconsin are limited to places with pack-area mean road densities of 0.7 mi/1 mi² (1.1 km/1 km²) or less (Mladenoff et al. 1995). Some studies have shown that a few large areas of low road density, even in a landscape of high average road density, may be the best indicator of suitable habitat for large vertebrates (Rudis 1995). Access and construction roads for the proposed action commonly are spurs from existing roads and range between 500 ft (152 m) and 1,000 ft (305 m) in length, which do not isolate or separate habitat patches. Furthermore, construction activities within montane woodlands, riparian corridors or major canyons will be minimal and widely distributed, resulting in negligible impacts to the composition or structure of Mexican gray wolf habitat.

Increased Legal and Unauthorized Access to Mexican Gray Wolf Habitat

Gray wolves experience negative interactions with humans and roads are a key facilitator (Thiel 1985). Increased human access to potential wolf habitat through the use of temporary proposed construction roads could reduce the quality of the habitat and human interactions may increase mortality (Mech 1973). The road closure techniques outlined in the SECTION 1.4 and the RA (URS 2003) will minimize unintended uses of these roads.

Accidental Wildfire

Increased road access may contribute to an increase in the frequency of human-caused ignitions in some areas (Gucinski et al. 2001). Because of their mobility, wolves will not likely be directly impacted by wildfires; however, these wildfires could potentially alter or destroy portions of prey species habitat. While the short-term effects of wildfires may affect prey species through loss of forage from the fire, increased herbaceous production in the years following a fire may improve habitat in the long term.

New roads also may act as firebreaks and improve response time of firefighters to wildfires, thereby preventing these fires from gaining in size and intensity. A study in southern California concluded that the road network had been a key factor in determining suppression strategies were used, both in firefighter access and because roads were

widely used for backfiring and burning-out operations (Salazar and Gonzalez-Caban 1987). Early studies of fuelbreak effectiveness in southern California came to similar conclusions (Green 1977). If deemed appropriate, new roads may allow fuelwood collection in areas currently not accessible, thereby reducing the density of downed, woody material, which is capable of carrying wildfires across the landscape. Fire prevention measures outlined in the Fire Prevention Plan will minimize the risks of wildfires associated with the proposed action.

Invasive Species

Roads may be the first point of entry for invasive species into a new landscape, and can serve as a corridor along which plants move farther into the landscape (Lonsdale and Lane 1994, Greenberg et al. 1997). Some invasive plants may then be able to move into adjacent patches of suitable habitat. Invasion by these plants may have significant biological and ecological effects if the species are able to disrupt the structure or function of an ecosystem. Roads constructed for the proposed action could allow the establishment or increased density of non-native plants, such as Lehmann's lovegrass, an invasive species that facilitates wildfires (McPherson 1995). Measures outlined in the Invasive Species Management Plan will minimize the introduction or spread of invasive species as a result of the proposed action.

2.7g Cumulative Effects

Cumulative effects include the effects of future state, local, or private actions that are reasonably certain to occur in the action area considered in this BA. While the action area for this species encompasses private, state, and federal lands, the habitat with the highest potential for occupancy by Mexican gray wolf occurs on USFS land in Santa Cruz County. Future federal actions will be subject to Section 7 consultation and will not be considered cumulative.

Although the amount of future private development within Santa Cruz County is unknown, many rural areas of Arizona are experiencing substantial growth. Between 1990 and 2000, Santa Cruz County grew 29.3 percent (U.S. Census Bureau 2000). Despite its distance from the proposed action, an increase in population in Nogales, Arizona and other regional population centers translates into an increased demand for recreational use of USFS land.

An undetermined level of border crossings by UDI also occurs within the action area and results in habitat damage from new roads, discarded trash, illegal campfires, and disturbance near water sources. These border crossings are likely to continue or increase.

2.7h Effects Determination and Incidental Take

Construction noise and activity associated with the proposed action may affect the Mexican gray wolf, but it is not likely to adversely affect the species because any disturbance will be widely distributed and short term in duration. Because the proposed action is not likely to adversely affect the Mexican gray wolf, no take is anticipated.

3.0 USFS SENSITIVE SPECIES

USFS Special status species are plant and wildlife species that are of concern because their populations are declining in size. In a letter dated 2 May 2002, AGFD listed 21 USFS Sensitive species that are known to occur within 3 mi (4.8 km) of the proposed corridor or may be expected to occur along the corridor if suitable habitat exists. The information listed in the letter was based on the AGFD Heritage Data Management System. In addition, 21 USFS Sensitive species known to occur within 5 mi (8 km) to 10 mi (16 km) of the proposed corridor have been included (AGFD letter dated 25 April 2002). AGFD species abstracts and other literature were reviewed for species' historical ranges and habitat preferences and field reconnaissance surveys were conducted along the entire corridor. However, species-specific surveys were impractical because of ongoing drought conditions in the project area, therefore the potential presence of sensitive species was assumed in all areas containing potential habitat. The 42 USFS Sensitive species that may occur on or near the proposed Central Corridor are listed in Table 3.

TABLE 3. SUMMARY OF EFFECTS ON U. S. FOREST SERVICE SENSITIVE SPECIES.

COMMON NAME <i>Scientific Name</i>	EFFECTS DETERMINATION	JUSTIFICATION
Alamos Deer Vetch <i>Lotus alamosanus</i>	May impact individuals of this species, but is not likely to result in a trend toward federal listing or loss of viability.	<ul style="list-style-type: none"> Minimal impacts to riparian habitat. Only small percentage of total population within project area may be impacted. Other viable populations occur outside of project area.
Arid Throne Fleabane <i>Erigeron arisolis</i>	May impact individuals of this species, but is not likely to result in a trend toward federal listing or loss of viability.	<ul style="list-style-type: none"> Only small percentage of total population within project area may be impacted. Other viable populations occur outside of project area.
Arizona Giant Sedge <i>Carex ultra</i>	May impact individuals of this species, but is not likely to result in a trend toward federal listing or loss of viability.	<ul style="list-style-type: none"> Minimal impacts to riparian habitat. Only small percentage of total population within project area may be impacted. Other viable populations occur outside of project area.
Arizona Metalmark <i>Calephelis rawsoni arizonensis</i>	May impact individuals of this species, but is not likely to result in a trend toward federal listing or loss of viability.	<ul style="list-style-type: none"> Only small percentage of total population within project area may be impacted. Populations of this species occur in isolated mountain ranges throughout southern Arizona. Mitigation plantings of host species will reduce impacts.
American Peregrine Falcon <i>Falco peregrinus anatum</i>	No Impacts.	<ul style="list-style-type: none"> Known occurrences and potential habitat are outside project area.
Bartram's Stonecrop <i>Graptopetalum bartramii</i>	No Impacts.	<ul style="list-style-type: none"> Known populations occur outside project area.
Beardless Chinch Weed <i>Pectis imberbis</i>	May impact individuals of this species, but is not likely to result in a trend toward federal listing or loss of viability.	<ul style="list-style-type: none"> Only small percentage of total population within project area may be impacted. Populations of this species occur in isolated mountain ranges throughout southern Arizona. Species is adapted to disturbances.
Broadleaf ground cherry <i>Physalis latiphysa</i>	No Impacts.	<ul style="list-style-type: none"> Known populations occur outside project area.

TABLE 3 (CONTINUED). SUMMARY OF EFFECTS ON U. S. FOREST SERVICE SENSITIVE SPECIES.		
COMMON NAME <i>Scientific Name</i>	EFFECTS DETERMINATION	JUSTIFICATION
Catalina Beardtongue <i>Penstemon discolor</i>	No Impacts.	<ul style="list-style-type: none"> Known populations occur outside project area.
Cave Myotis <i>Myotis velifer</i>	May impact individuals of this species, but is not likely to result in a trend toward federal listing or loss of viability.	<ul style="list-style-type: none"> No known roosts within project area. Only small percentage of foraging habitat within project area may be impacted. Populations of this species occur throughout southern Arizona.
Chiltepine <i>Capsicum annuum</i> var. <i>glabriusculum</i>	May impact individuals of this species, but is not likely to result in a trend toward federal listing or loss of viability.	<ul style="list-style-type: none"> Only small percentage of total population within project area may be impacted. Populations of this species occur in isolated mountain ranges throughout southern Arizona.
Chihuahuan Sedge <i>Carex chihuahuensis</i>	No Impacts.	<ul style="list-style-type: none"> Known populations occur outside project area.
Chiricahua Mountain Brookweed <i>Samolus vagans</i>	No Impacts.	<ul style="list-style-type: none"> No construction in perennial aquatic habitats.
Five-Stripped Sparrow <i>Aimophila quinquestriata</i>	No Impacts.	<ul style="list-style-type: none"> Potential habitat and know occurrences are outside project area.
Foetid Passionflower <i>Passiflora foetida</i>	No Impacts.	<ul style="list-style-type: none"> Known populations occur outside project area.
Gentry Indigo Bush <i>Dalea tentaculoides</i> Giant Spotted Whiptail <i>Cnemidophorus burti</i> <i>strictogrammus</i>	No Impacts.	<ul style="list-style-type: none"> Known populations occur outside project area. Known populations occur outside project area.
Large-Flowered Blue Star <i>Amsonia grandiflora</i>	May impact individuals of this species, but is not likely to result in a trend toward federal listing or loss of viability.	<ul style="list-style-type: none"> Only small percentage of total population within project area may be impacted. Other viable populations occur outside of project area.

TABLE 3 (CONTINUED). SUMMARY OF EFFECTS ON U. S. FOREST SERVICE SENSITIVE SPECIES.		
COMMON NAME <i>Scientific Name</i>	EFFECTS DETERMINATION	JUSTIFICATION
Lowland Leopard Frog <i>Rana yavapaiensis</i>	No Impacts.	<ul style="list-style-type: none"> Known populations occur outside project area. No construction in perennial aquatic habitats.
Lumholtz Nightshade <i>Solanum lumholtzianum</i>	May impact individuals of this species, but is not likely to result in a trend toward federal listing or loss of viability.	<ul style="list-style-type: none"> Minimal impacts to riparian habitat. Only small percentage of total population within project area may be impacted. Populations of this species occur in isolated mountain ranges throughout southern Arizona.
Mexican Garter Snake <i>Thamnophis eques megalops</i>	No Impacts.	<ul style="list-style-type: none"> No construction in perennial aquatic habitats. Minimal impacts to riparian habitat.
Mock-Pennyroyal <i>Hedeoma dentatum</i>	May impact individuals of this species, but is not likely to result in a trend toward federal listing or loss of viability.	<ul style="list-style-type: none"> Only small percentage of total population within project area may be impacted. Populations of this species occur in isolated mountain ranges throughout southern Arizona.
Nodding Blue-eyed Grass <i>Sisyrinchium cernuum</i>	No Impacts.	<ul style="list-style-type: none"> Known populations occur outside project area.
Northern Gray Hawk <i>Asturina nitida maxima</i>	May impact individuals of this species, but is not likely to result in a trend toward federal listing or loss of viability.	<ul style="list-style-type: none"> Distance of habitat from project area will attenuate effects. Only small percentage of total population within project area may be impacted.
Pima Indian mallow <i>Abutilon parishii</i>	No Impacts.	<ul style="list-style-type: none"> Known populations occur outside project area.
Santa Cruz Beehive Cactus <i>Coryphantha recurvata</i>	May impact individuals of this species, but is not likely to result in a trend toward federal listing or loss of viability.	<ul style="list-style-type: none"> Only small percentage of total population within project area may be impacted. Other viable populations occur outside of project area.
Santa Cruz Star Leaf <i>Choisya mollis</i>	No Impacts.	<ul style="list-style-type: none"> Known populations occur outside project area.
Santa Cruz Striped Agave <i>Agave parviflora</i> ssp. <i>parviflora</i>	May impact individuals of this species, but is not likely to result in a trend toward federal listing or loss of viability.	<ul style="list-style-type: none"> Only small percentage of total population within project area may be impacted. Plants occur throughout Nogales Ranger District. Mitigation plantings of agave will reduce impacts.

TABLE 3 (CONTINUED). SUMMARY OF EFFECTS ON U. S. FOREST SERVICE SENSITIVE SPECIES.		
COMMON NAME <i>Scientific Name</i>	EFFECTS DETERMINATION	JUSTIFICATION
Seeman Groundsel <i>Senecio carlomasonii</i>	No Impacts.	<ul style="list-style-type: none"> Known populations occur outside project area.
Sonoran Noseburn <i>Tragia laciniata</i>	May impact individuals of this species, but is not likely to result in a trend toward federal listing or loss of viability.	<ul style="list-style-type: none"> Only small percentage of total population within project area may be impacted. Populations of this species occur in isolated mountain ranges throughout southern Arizona.
Southern Pocket Gopher <i>Thomomys umbrinus intermedius</i>	May impact individuals of this species, but is not likely to result in a trend toward federal listing or loss of viability.	<ul style="list-style-type: none"> Only small percentage of total population within project area may be impacted. Other viable populations occur outside of project area.
Superb Beardtongue <i>Penstemon superbus</i>	May impact individuals of this species, but is not likely to result in a trend toward federal listing or loss of viability.	<ul style="list-style-type: none"> Only small percentage of total population within project area may be impacted. Populations of this species occur in isolated mountain ranges throughout southern Arizona.
Supine Bean <i>Macroptilium supinum</i>	May impact individuals of this species, but is not likely to result in a trend toward federal listing or loss of viability.	<ul style="list-style-type: none"> Pre-construction surveys will be conducted and, if necessary, mitigation measures will be coordinated with USFS personnel.
Sweet Acacia <i>Acacia smallii</i>	May impact individuals of this species, but is not likely to result in a trend toward federal listing or loss of viability.	<ul style="list-style-type: none"> Only small percentage of total population within project area may be impacted. Other viable populations occur outside of project area.
Thurber Hoary Pea <i>Tephrosia thurberi</i>	May impact individuals of this species, but is not likely to result in a trend toward federal listing or loss of viability.	<ul style="list-style-type: none"> Only small percentage of total population within project area may be impacted. Populations of this species occur in isolated mountain ranges throughout southern Arizona.
Thurber's Morning-glory <i>Ipomoea thurberi</i>	May impact individuals of this species, but is not likely to result in a trend toward federal listing or loss of viability.	<ul style="list-style-type: none"> Only small percentage of total population within project area may be impacted. Populations of this species occur in isolated mountain ranges throughout southern Arizona.
Virlet Paspalum <i>Paspalum virletti</i>	No Impacts.	<ul style="list-style-type: none"> Known populations occur outside project area.

TABLE 3 (CONTINUED). SUMMARY OF EFFECTS ON U. S. FOREST SERVICE SENSITIVE SPECIES.		
COMMON NAME <i>Scientific Name</i>	EFFECTS DETERMINATION	JUSTIFICATION
Weeping Muhly <i>Muhlenbergia xerophila</i>	May impact individuals of this species, but is not likely to result in a trend toward federal listing or loss of viability.	<ul style="list-style-type: none"> • Only small percentage of total population within project area may be impacted. • Populations of this species occur in isolated mountain ranges throughout southern Arizona.
Western Barking Frog <i>Eleutherodactylus augusti cactorum</i>	No Impacts.	<ul style="list-style-type: none"> • Known populations occur outside project area.
Western Yellow-billed Cuckoo <i>Coccyzus americanus</i>	May impact individuals of this species, but is not likely to result in a trend toward federal listing or loss of viability.	<ul style="list-style-type: none"> • Minimal impacts to riparian habitat. • Only small percentage of total population within project area may be impacted. • Other viable populations occur outside of project area.
Wiggins Milkweed Vine <i>Metastelma mexicanum</i>	May impact individuals of this species, but is not likely to result in a trend toward federal listing or loss of viability.	<ul style="list-style-type: none"> • Populations within Arizona appear stable. • Only small percentage of total population within project area may be impacted. • Populations of this species occur in isolated mountain ranges throughout southern Arizona.
Wooly Fleabane <i>Laennecia eriophylla</i>	No Impacts.	<ul style="list-style-type: none"> • Potential habitat and know occurrences are outside project area.

3.1 PLANTS

Alamos deer vetch (*Lotus alamosanus*)

Alamos deer vetch is a perennial herb found in southern Arizona, and Sonora, Chihuahua, and Durango, Mexico. Within Arizona, this plant is found in Sycamore Canyon and the Pajarito Mountains of Santa Cruz County, and near Garden Valley in Maricopa County. This plant is considered a wetland obligate species that is restricted to stream banks in canyons at elevations ranging from 3,500 ft (1,067 m) to 5,500 ft (1,676 m) (AGFD 1999a). Within the Nogales RD, this plant occurs in the Sycamore Canyon and Peña Blanca Canyon areas (T. Newman, CNF, pers. comm., 20 August 2002).

Population trends for Alamos deer vetch are unknown (AGFD 1999a). The proposed transmission line may cross potential Alamos deer vetch habitat; however, construction within riparian habitats will be minimized to the greatest extent possible. Furthermore, viable populations occur outside of the project area, including the Gooding RNA. There may be an impact to individual plants during development of the line; however, disturbance will be limited to a few individuals and is not likely to result in a trend toward federal listing or loss of viability.

Arid throne fleabane (*Erigeron arisolis*)

Arid throne fleabane is an annual to short-lived perennial forb that occurs in Arizona, southwestern New Mexico and Sonora, Mexico. Within Arizona, this plant is found in Apache, Cochise, Pima, and Santa Cruz counties. This species is typically found on moist rocky soils in grasslands, grassy openings within oak woodlands, and roadsides at elevations between 4,200 ft (1,280 m) and 5,500 ft (1,676 m) (AGFD 2000a). On the CNF Nogales RD, it has been documented from Box Canyon and Ruby Roads (T. Newman, CNF, pers. comm., 20 August 2002).

Arid throne fleabane favors moist areas in grasslands and grassy openings in oak woodlands, areas also favored by livestock for grazing (AGFD 2000a). The proposed transmission line parallels Ruby Road, a known location for this species. Placement of the transmission line may impact individual arid throne fleabane, however because of the linear nature of the project, only a small percentage of the population within the project area may be impacted. Furthermore, populations of this species occur outside of the project area. Therefore, impacts are not likely to result in a trend toward federal listing or loss of viability.

Arizona giant sedge (*Carex ultra*)

Arizona giant sedge is the largest sedge found in Arizona. Its range includes southeast Arizona, extreme southwest New Mexico (Hidalgo County, Indian Springs in the Pelocillos) and Mexico (Sonora and Coahila). Within Arizona, this sedge is found in Cochise, Graham, Pinal, Yavapai, Pima (Santa Rita Mountains and the Rincon Valley), and Santa Cruz counties (Santa Rita and Atascosa mountains). Typically only 1 patch per mountain has been found. Like other sedges, this plant is associated with moist soil

near perennial wet springs and streams and undulating rocky-gravelly terrain at elevations ranging from 2,040 ft (622 m) to 6,000 ft (1,829 m) (AGFD 2000b). Within the Nogales RD, Arizona giant sedge is found in Sycamore Canyon and Mule Ridge in the Atascosa Mountains, and at Deering Spring and Big Casa Blanca Canyon in the Santa Rita Mountains (T. Newman, CNF, pers. comm., 20 August 2002).

Small populations of this sedge are vulnerable to local disturbance of aquatic or riparian habitat (AGFD 2000b). The proposed transmission line may cross potential Arizona giant sedge habitat; however, no construction will occur in perennial aquatic habitats and construction within riparian habitats will be minimized to the greatest extent possible. There may be an impact to individual plants during development of the line; however because of the linear nature of the project, only a small percentage of the population within the project area may be impacted. Therefore, impacts are not likely to result in a trend toward federal listing or loss of viability.

Bartram's stonecrop (*Graptopetalum bartramii*)

Bartram's stonecrop is a small succulent perennial found in southern Arizona and Chihuahua, Mexico (one record). In Arizona, this plant occurs in Santa Cruz County within the Patagonia, Santa Rita, and Tumacacori Mountains, in Pima County within the Baboquivari, Dragoon, and Rincon mountains, and in Cochise County within the Chiricahua Mountains. Habitat for Bartram's stonecrop consists of cracks in rocky outcrops within shrub live oak-grassland communities located on the sides of rugged canyons. This plant is usually found in heavy litter cover and shade where moisture drips from rocks at elevations ranging from 3,900 ft (1,189 m) to 6,700 ft (2,042 m) (AGFD 1997a). Bartram's stonecrop plants are found on the west side of the Nogales RD in Tres Amigos Gulch; Sycamore, Peña Blanca, Alamo, and Peñasco canyons; in the vicinity of Montana Peak and Peña Blanca Lake (T. Newman, CNF, pers. comm., 20 August 2002).

Bartram's stonecrop populations are typically small and isolated. Illegal collection of the plant is the main management issue at this time. Other factors that may affect populations include mining and mineral exploration, habitat alteration due to livestock grazing, trampling by cattle and recreationists, and road construction and maintenance. Bartram's stonecrop populations are typically small and isolated but illegal collection of the plant is the main management issue at this time. Other factors that may affect populations include mining and mineral exploration, habitat alteration due to livestock grazing, trampling by cattle and recreationists, and road construction and maintenance (AGFD 1997a). The proposed TEP transmission line does not cross known Bartram's stonecrop populations within the Nogales RD, therefore placement of the transmission line will not impact this species.

Beardless chinch weed (*Pectis imberbis*)

Beardless chinch weed is a perennial herb that is found in southern Arizona, western Chihuahua and eastern Sonora, Mexico. Within Arizona, this plant can be found in Cochise, Pima, and Santa Cruz counties (within Santa Cruz County it is found along Ruby Road in the Atascosa Mountains and in the Red Rock area of Canelo Hills). Habitat for this species consists of open areas in grassland and oak-grassland

communities. Beardless chinch weed has an extremely broad habitat range and can be found at elevations from 4,000 ft (1,219 m) to 5,000 ft (1,524 m) (AGFD 1998a).

Populations of beardless chinch weed may be susceptible to impacts from grazing and road maintenance activities but the species is adapted to disturbances and grows along road cuts (AGFD 1998a). The proposed transmission line crosses over known beardless chinch weed populations within the Nogales RD. Placement of the transmission line may impact individual beardless chinch weed, however because of the linear nature of the project, only a small percentage of the population within the project area may be impacted. Furthermore, populations of this species occur in isolated mountain ranges throughout southern Arizona. Therefore, impacts to beardless chinch weed are not likely to result in a trend toward federal listing or loss of viability.

Broadleaf ground cherry (*Physalis latiphysa*)

Broadleaf ground cherry is an herbaceous annual found in southern Arizona. This plant can be found in the San Bernardino Valley of Cochise County, the Pinaleno Mountains of Graham County, in the vicinity of Arivaca Creek in Pima County, and the Santa Cruz River of Santa Cruz County. Habitat for the broad-leaf ground cherry consists of washes, often in the shade of shrubs and boulders, desertscrub vegetation, and grasslands at elevations ranging from 914 to 1,372 m (3,000 – 4,500 feet) (AGFD 2000c). There are no known sites for this plant in the Nogales RD. The nearest locations are northwest of Arivaca Lake and in the vicinity of Tubac on the Santa Cruz River (T. Newman, CNF, pers. comm., 20 August 2002).

There is no information on the potential effects of land use activities, such as utility placement, on the population status of broad-leaf ground cherry (AGFD 2000c). The proposed TEP transmission line does not cross known broadleaf ground cherry populations within the Nogales RD, therefore placement of the transmission line will not impact this species.

Catalina beardtongue (*Penstemon discolor*)

Catalina beardtongue is a perennial herbaceous sub-shrub found in southern Arizona. This shrub is found in Cochise, Graham, Pinal, Pima (within the Santa Catalina Mountains), and Santa Cruz counties (within the Atascosa and Tumacacori mountains). Habitat for Catalina beardtongue consists of bare rock outcrops, barren soil outcrops, and bedrock openings in chaparral or pine-oak woodlands at elevations ranging from 4,120 ft (1,256 m) to 7,600 ft (2,316) (AGFD 1999b). On the Nogales RD, this shrub occurs in the upper end of Peck Canyon, Corral Nuevo, and the adjacent Bartalo Mountain (Cedar Canyon), typically on whitish volcanic ash (T. Newman, CNF, pers. comm., 20 August 2002).

Rock climbers threaten some populations of this plant but few other threats exist (AGFD 1999b). The proposed TEP transmission line does not cross known Catalina beardtongue populations within the Nogales RD, therefore placement of the transmission line will not impact this species.

Chiltepine (*Capsicum annuum* var. *glabriusculum*)

Chiltepine is an herbaceous to woody perennial shrub that is found in south Texas, southern New Mexico, southern Arizona, and south to tropical America. Within Arizona, a few populations of this plant are found in the Chiricahua, Tumacacori, Baboquivari, and Ajo Mountains. This plant occurs in protected, frost-free canyons in oak woodlands of slopes at less than 4,500 ft (1,372 m) elevation (typically found at elevations ranging from 3,600 ft [1,097 m] to 4,400 ft [1,341 m]). Chiltepine plants grow under nurse shrubs and usually are associated with rock ledges and outcrops. Within the Nogales RD, there are populations in the Tumacacori Mountains and Cobre Ridge area, and there are suspected populations on the west side of the RD (AGFD 1991a; T. Newman, CNF, pers. comm., 20 August 2002).

This plant is declining in some areas because of drought, overgrazing, and local over-collection of berries (AGFD 1991a). Placement of the transmission line may impact individual chiltepine plants, however because of the linear nature of the project, only a small percentage of the population within the project area may be impacted. Furthermore, populations of this species occur in isolated mountain ranges throughout southern Arizona. Therefore, impacts to chiltepine are not likely to result in a trend toward federal listing or loss of viability.

Chihuahuan sedge (*Carex chihuahuensis*)

Chihuahuan sedge is a grass-like perennial plant that occurs in southeastern Arizona, New Mexico (Hidalgo County), and Mexico (Sonora and Chihuahua). Within Arizona, this plant ranges from Cochise, Graham, Gila, Pima (Santa Catalina, San Luis, and Rincon mountains), and Santa Cruz counties (Atascosa and Santa Rita mountains, and the Santa Cruz River). Chihuahuan sedge can be found in wet soils along streambeds and in shallower draws of pine-oak forests and riparian woodlands. It also is found in wet meadows, cienegas, marshy areas, and canyon bottoms from 1,100 ft (335 m) to 8,000 ft (AGFD 1999c). Within the Nogales RD, this plant has been found near Arivaca Lake (on private land), Sycamore Canyon, and south of Bear Valley (T. Newman, CNF, pers. comm., 20 August 2002).

There is no information on the potential effects of land use activities, such as utility placement on the population status of Chihuahuan sedge (AGFD 1999c). The proposed transmission line may cross potential Chihuahuan sedge habitat; however, construction within riparian habitats will be minimized to the greatest extent possible. There may be an impact to individual plants during development of the line; however because of the linear nature of the project, only a small percentage of the population within the project area may be impacted. Therefore, impacts are not likely to result in a trend toward federal listing or loss of viability.

Chiricahua Mountain brookweed (*Samolus vagans*)

Chiricahua Mountain brookweed is a perennial herb found in southeastern Arizona, western Chihuahua, and eastern Sonora, Mexico. This plant apparently reaches its southern limit in southern Sonora, Mexico. Within Arizona, this species is found in the Huachuca Mountains of Cochise County, the Rincon, Santa Catalina, and Santa Rita

mountains of Pima County, and the Canelo Hills and Pajarito mountains of Santa Cruz County. The Chiricahua Mountain brookweed is confined to areas with permanent water, such as springs, seeps, and in and along streams at elevations ranging from 1,219 to 2,195 m (4,000 – 7,200 ft) (AGFD 1999d). Within the Nogales RD, this plant occurs in Florida Canyon of the Santa Rita Mountains and in Sycamore Canyon of the Atascosa Mountains (T. Newman, CNF, pers. comm., 20 August 2002).

There is no information on the potential effects of land use activities, such as utility placement, on the population status of Chiricahua Mountain brookweed (AGFD 1999d). Because no construction will occur within perennial aquatic habitats, the proposed action will have no effect on the population status of the Chiricahua Mountain brookweed.

Foetid passionflower (*Passiflora foetida*)

The foetid passionflower is a herbaceous vine found in southeastern Texas and the Rio Grande Valley, southern Arizona, and southward throughout Mexico, Central and South America, and the West Indies. Within Arizona, this species is found in the Baboquivari Mountains, Arivaca, and Las Guijas Mountains of Pima County and in California Gulch and the Bartlett Mountains of Santa Cruz County. In Arizona, this plant occurs on hillsides and canyons of the Lower Sonoran zone from 1,067 to 1,707 m (3,500 – 5,600 ft) in elevation (AGFD 2000c). Within the Nogales RD, foetid passionflowers have been recorded in the California Gulch and Holden Canyon areas (T. Newman, CNF, pers. comm., 20 August 2002).

There is no information on the potential effects of land use activities, such as utility placement, on the population status of foetid passionflower (AGFD 2000c). Known locations of this plant occur outside of the proposed TEP transmission line corridor, therefore the proposed TEP transmission line will have no effect on the population status of the foetid passionflower

Gentry indigo bush (*Dalea tentaculoides*)

The Gentry indigo bush is an herbaceous perennial shrub found primarily in southern Arizona, but its range may extend into Mexico. Within Arizona, this shrub is found in the Sycamore Canyon drainage of the Atascosa Mountains, in the Pajarito Mountains of Santa Cruz County, and within the Baboquivari Mountains (1930s record) and Mendoza Canyon (1965 record) of Pima County. Gentry indigo bush is typically found along canyon bottoms on cobble terraces subject to occasional flooding and seems to prefer disturbance-prone environments at elevations ranging from 1,097 to 1,341 m (3,600 – 4,400 ft) (AGFD 1998b). Historic collection records indicate that this plant may grow on rocky hillsides. Within the Nogales RD, this plant has been recorded in Sycamore Canyon, in the vicinity of Peñasco Canyon, Kaiser Canyon, and north of Manzanita Mountain (T. Newman, CNF, pers. comm., 20 August 2002).

Potential threats to Gentry indigo bush populations are cattle grazing, recreational foot traffic, and flooding events that eliminate terraces occupied by this species (AGFD 1998b). Known locations of this plant occur outside of the proposed TEP transmission line corridor, therefore the proposed TEP transmission line will have no effect on the population status of the Gentry indigo bush.

Large-flowered blue star (*Amsonia grandiflora*)

The large-flowered blue star is an herbaceous perennial that is found in northern Sonora and Durango, Mexico, and southern Arizona. Within Arizona, this plant is found in the Patagonia, Atascosa/Pajarito mountains of Santa Cruz and Pima counties. Habitat for this species consists of canyon bottoms in oak woodlands typically dominated by Emory oak and Mexican blue oak; however, site-specific qualities are inconsistent. Large-flowered blue star plants have adapted to rock fall disturbance and are typically found at elevations ranging from 1,189 to 1,372 m (3,900 to 4,500 ft) (AGFD 1998c). Within the west side of the Nogales RD, this plant occurs at Peña Blanca and Arivaca Lakes, Sycamore Canyon, Chiminea Canyon, California Gulch, and near Ruby (T. Newman, CNF, pers. comm., 20 August 2002).

Populations of large-flowered blue star are rare, with only 15 to 20 populations within 2 mountain ranges as the total world distribution, but populations seem to be stable. This plant is highly susceptible to disturbance, and expanding development in the Nogales area (AGFD 1998c) may impact populations. The proposed TEP transmission line crosses near a known large-flowered blue star population in Peña Blanca Canyon, and some individual plants, comprising a small percentage of the total population, may be impacted. Therefore, impacts are not likely to result in a trend toward federal listing or loss of viability.

Lumholtz nightshade (*Solanum lumholtzianum*)

The Lumholtz nightshade is an herbaceous annual that is found in southern Arizona and northern Mexico. Within Arizona, this plant is found in the Arivaca and San Luis Mountains of Pima County and the Patagonia, Atascosa, and Santa Rita Mountains of Santa Cruz County. Lumholtz nightshade plants are typically found in washes and low ground near wet depressions and along stream banks from 914 to 1,402 m (3,000 – 4,600 ft) elevation in desert grassland plant communities. This plant is also often found in disturbed, weedy areas (AGFD 2000d). Within the Nogales RD, this nightshade is found in the vicinity of Arivaca, Ruby, California Gulch, Nogales, Cobre Ridge, and Oro Blanco Wash (T. Newman, CNF, pers. comm., 20 August 2002).

There is no information on the potential effects of land use activities, such as utility placement, on the population status of Lumholtz nightshade (AGFD 2000d). The proposed transmission line may cross potential habitat for this species; however, construction within riparian habitats will be minimized to the greatest extent possible. Placement of the transmission line may impact individual plants, however because of the linear nature of the project, only a small percentage of the population within the project area may be impacted. Furthermore, populations of this species occur in isolated mountain ranges throughout southern Arizona. Therefore, impacts to this species are not likely to result in a trend toward federal listing or loss of viability.

Mock-pennyroyal (*Hedeoma dentatum*)

The mock-pennyroyal is an herbaceous perennial plant found in southeastern Arizona and northern Sonora, Mexico. Within Arizona, this plant is found in the Chiricahua, Huachuca, Mule, Whetstone, and Winchester mountains of Cochise County, the Pinaleno

Mountains of Graham County, the Baboquivari, Rincon, and Santa Cruz mountains of Pima County, and the Atascosa, Mustang, Pajarito, and Santa Rita mountains of Santa Cruz County. Habitat for this plant consists of oak woodland, oak-pine forest, and pine forest. It can be found on open roadcuts, steep rocky outcrops, and gravelly slopes in wooded canyons with open to full sunlight at elevations ranging from 1,173 to 2,500 m (3,850 – 8,200 ft) (AGFD 2000e).

Populations of mock-pennyroyal seem to be restricted to a relatively small geographic area, and populations are apparently small. Because habitat for this species is widespread, placement of the transmission line may impact individual plants. However because of the linear nature of the project, only a small percentage of the population within the project area may be impacted. Furthermore, populations of this species occur in isolated mountain ranges throughout southern Arizona. Therefore, impacts to this species are not likely to result in a trend toward federal listing or loss of viability.

Nodding blue-eyed grass (*Sisyrinchium cernuum*)

Nodding blue-eyed grass is a perennial forb with grass-like leaves that occurs in southeastern Arizona, west Texas, and Mexico. Within Pima and Santa Cruz counties, Arizona it occurs in the Pajarito, Santa Rita, Atascosa, and Rincon mountains as well as Sycamore Canyon. This species can be found in desert grassland and pine-oak woodlands from 1,006 to 2,438 m (3,300 – 8,000 ft) in elevation along streams in partial shade and in canyon bottoms. It grows in wet soil by seeps, pools, or springs in desert scrub. It has also been found on sandy stream banks. On the Nogales RD, this plant has been found at 1,189 m (3,900 ft) in Sycamore Canyon on the west side and at 1,402 m (4,600 ft) in Big Casa Blanca Canyon in the Santa Rita Mountains (AGFD 1999e). The known location of this plant in Sycamore Canyon is within the Goodding RNA, located approximately 1.6 km (1 mi) west of the proposed ROW (T. Newman, CNF, pers. comm., 20 August 2002).

There is no information on the potential effects of land use activities, such as utility placement, on the population status of nodding blue-eyed grass (AGFD 1999e). However, this species is not likely to be affected by the proposed placement of a transmission line within the Nogales RD. The proposed transmission line will not cross over or near known locations of this plant within the Goodding RNA. Therefore, placement of the TEP transmission line from Sahuarita to Nogales will have no impact on the nodding blue-eyed grass.

Pima Indian mallow (*Abutilon parishii*)

The Pima Indian mallow is a perennial woody-based plant with herbaceous branches. This plant is known from 84 populations in 17 mountain ranges from near the town of Bagdad in central Arizona to Nachopouli Canyon, Sonora, Mexico. Within Arizona, Pima Indian mallow are found in the Superstition Mountains of Maricopa County, the Santa Catalina, Rincon, Silverbell, and Tucson mountains of Pima County, the Mineral Hills, Superstition, Picacho, Tortolito, and Dripping Springs mountains of Pinal County, the Santa Rita and Tumacacori mountains of Santa Cruz County, and the Little Shipp Wash and Cottonwood Creek areas near Bagdad in Yavapai County. This plant has also

been identified within Sabino Canyon in Pima County. Pima Indian mallow are typically found in mesic situations in full sun within higher elevations of Sonoran desertscrub. They can be found on rocky slopes, cliff bases, lower side slopes and ledges of canyons among rocks and boulders. In riparian zones, this plant occurs on flat secondary terraces but typically not in canyon bottoms. Pima Indian mallow are often found near trails, probably because of the trails influence on the light, heat, and water on the micro-habitat. This species is found at elevations ranging from 900 to 1,440 m (3,000 to 4,800 feet) (AGFD 1997b). Within the Nogales RD, this plant occurs in the Devils Cash Box area of the Santa Rita Mountains and within Peck Canyon of the Tumacacori Mountains (T. Newman, CNF, pers. comm., 20 August 2002).

In Arizona, few threats exist to the populations of Pima Indian mallow because this plant grows in steep areas eliminating grazing pressures, and neither light fires nor freezing temperatures cause harm to it (AGFD 1997b). The proposed transmission line will not cross over or near known locations of this plant; therefore, placement of the TEP transmission line will have no impact on the population status of the Pima Indian mallow.

Santa Cruz beehive cactus (*Coryphantha recurvata*)

The Santa Cruz beehive cactus is a succulent perennial that occurs in southern Arizona and northern Sonora (about 20 km [12.4 mi] south of the international border), Mexico. Within Arizona, this species occurs in western Santa Cruz County from Nogales and the Tumacacori Mountains west to the Atascosa/Pajarito mountains. Santa Cruz beehive cacti are found in alluvial soils of valleys and foothills in grassland and oak woodland habitats from 1,219 to 1,829 m (4,000 – 6,000 ft). These plants are either on rocky hillsides with high grass cover or in rock crevices where runoff accumulates and provides a more favorable moisture relationship than the surrounding soils (AGFD 1998d). Within the Nogales RD known plant locations have increased since 1997 (813 plant clumps in 1997, 807 plant clumps in 1998, and 175 in 1999) (T. Newman, CNF, pers. comm., 20 August 2002).

Accessible populations of the Santa Cruz beehive cactus have declined due to collection, but the status of populations beyond accessible areas is unknown (AGFD 1998d). The proposed TEP transmission line crosses over several known Santa Cruz beehive cactus populations within the Nogales RD. Placement of the transmission line may impact individual plants, however because of the linear nature of the project, only a small percentage of the population within the project area may be impacted. Furthermore, populations of this species occur outside of the project area. Therefore, impacts are not likely to result in a trend toward federal listing or loss of viability.

Santa Cruz star leaf (*Choisya mollis*)

The Santa Cruz star leaf is a perennial shrub that occurs in southern Arizona within the Atascosa, Pajarito, and Tumacacori mountains of Santa Cruz County. Santa Cruz star leaf plants are found primarily within madrean evergreen woodland communities from 1,067 to 1,524 m (3,500 – 5,000 ft) in elevation. This plant is usually found in canyon bottoms and slopes, usually in the shade of oaks and other trees, or rock outcrops (AGFD

1999f). Santa Cruz star leaf plants have been found throughout the eastern portion of the Nogales RD (T. Newman, CNF, pers. comm., 20 August 2002).

Santa Cruz star leaf are typically found in rugged and remote mountainous areas where human activity is low and the likelihood of disturbance or removal of plants is minimal. However, the species population trend is unknown and existing populations are relatively rare, have a restricted range, and are only found within specific habitats (AGFD 1999f). The proposed transmission line will not cross over or near known locations of this plant; therefore, placement of the TEP transmission line will have no impact on the Santa Cruz star leaf.

Santa Cruz striped agave (*Agave parviflora* ssp. *parviflora*)

Santa Cruz striped agave is a small perennial succulent found in southern Arizona and northern Mexico. Within Arizona, this species is found near Arivaca in Pima County, and in the Las Guijas, Pajarito, Patagonia, Santa Rita, and Atascosa mountains of Santa Cruz County. Habitat for this agave consists of rocky or gravelly slopes of middle elevation mountains, in desert grassland or oak woodlands. This plant appears to prefer soils on rounded ridge-tops where grasses and shrubs are sparse and soil is bare or nearly so (AGFD 1998e). Santa Cruz striped agave have been found throughout the Nogales RD (primarily within the Atascosa, Pajarito, San Luis, and Las Guijas mountains), and in recent years the documented number of individual plants and number of locations has increased for this area (T. Newman, CNF, pers. comm., 20 August 2002).

Some populations of Santa Cruz striped agave have declined due to illegal collection and loss of habitat due to mining and road construction. Livestock grazing has caused degradation of habitat and browsing of flower stalks (AGFD 1998e). The proposed TEP transmission line crosses areas with known populations of Santa Cruz striped agave and there may be an impact to individual plants during development of the line. Placement of the transmission line may impact individual plants, however because of the linear nature of the project, only a small percentage of the population within the project area may be impacted. Furthermore, populations of this species occur outside of the project area and transplanting of agave plants in project area will minimize impacts. Therefore, impacts are not likely to result in a trend toward federal listing or loss of viability.

Seeman groundsel (*Senecio carlomasonii*)

The seeman groundsel is a perennial herb or subshrub found in southern Arizona and Mexico (Sonora, Chihuahua, Nayarit). Within Arizona, this plant is found in the Chiricahua and Huachuca mountains of Cochise County, the Baboquivari and Santa Rita mountains of Pima County, and the Santa Rita, Pajarito, and Peña Blanca mountains of Santa Cruz County (AGFD 2000f). Within the Nogales RD, seeman groundsel have been recorded in the Peña Blanca Lake and Sycamore Canyon areas (T. Newman, CNF, pers. comm., 20 August 2002).

There is no information on the potential effects of land use activities, such as utility placement, on the population status of seeman groundsel (AGFD 2000f). A potential threat to seeman groundsel habitat may be trampling by hikers. The proposed

transmission line will not cross over or near known locations of this plant; therefore, placement of the TEP transmission line will have no impact on the population status of the seaman groundsel.

Sonoran noseburn (*Tragia laciniata*)

Sonoran noseburn is an herbaceous perennial that occurs in southern Arizona, Mexico (Sonora and Chihuahua), and possibly New Mexico. Within Arizona this plant can be found in Cochise County in the Huachuca Mountains and Canelo Hills, in Pima County in the Santa Rita Mountains, and in Santa Cruz County in the Atascosa Mountains (Sycamore Canyon), Patagonia Mountains, Pajarito Mountains, Canelo Hills (O'Donnell Canyon), and Santa Rita Mountains. Sonoran noseburn typically occur at elevations of 1,067 to 1,722 m (3,500 – 5,650 ft) along streams and canyon bottoms, on shaded hillsides within the upper parts of the Lower Sonoran and Upper Sonoran biotic communities, and open woodland areas (AGFD 2000g). This species has been found in canyons, along streams, and near roadways of the Nogales RD (AGFD 2000g).

There is no information on the potential effects of land use activities, such as utility placement, on the population status of Sonoran noseburn (AGFD 2000g). Placement of the transmission line may impact individual plants, however because of the linear nature of the project, only a small percentage of the population within the project area may be impacted. Furthermore, populations of this species occur in isolated mountain ranges throughout southern Arizona. Therefore, impacts to this species are not likely to result in a trend toward federal listing or loss of viability.

Superb beardtongue (*Penstemon superbus*)

The superb beardtongue is a perennial herbaceous forb found in southeastern Arizona, New Mexico, and Mexico (Chihuahua). Within southern Arizona, this species is found in Pima County in the Santa Catalina and Santa Rita mountains, and in Santa Cruz County within the Tumacacori Mountains. This plant is generally found in rocky canyons, dry hillsides, and along washes in sandy or gravelly soils at elevations between 945 and 1,676 m (3,100 – 5,500 ft) (AGFD 2000h). Within the Nogales RD, it has been found in Rock Corral Canyon and Box Canyon (T. Newman, CNF, pers. comm., 20 August 2002).

There is no information on the potential effects of land use activities, such as utility placement, on the population status of superb beardtongue (AGFD 2000h). Placement of the transmission line may impact individual plants, however because of the linear nature of the project, only a small percentage of the population within the project area may be impacted. Furthermore, populations of this species occur in isolated mountain ranges throughout southern Arizona. Therefore, impacts to this species are not likely to result in a trend toward federal listing or loss of viability.

Supine bean (*Macroptilium supinum*)

The supine bean is a perennial herb that grows in colonies and produces underground fruits. The total range for this species includes Santa Cruz County, Arizona, south into Mexico, including the states of Sonoran and Nayarit. Within Arizona, this plant can be found in the Atascosa/Pajarito, San Luis, and Patagonia Mountains, and the southern

portion of the Santa Cruz River drainage in Santa Cruz County (much of this area is within the Nogales RD). Supine bean are typically found along ridge tops and gentle slopes of rolling hills in semi-desert grassland or grassy openings in oak-juniper woodlands at elevations between 1,097 and 1,494 m (3,600 – 4,900 ft) (AGFD 1999g).

There are currently an estimated 12 populations of this species in Arizona. Populations range from small (around 20 individuals) to relatively large (around 3,500 individuals). A 43% decline in a monitored population was recorded from 1989 to 1993. This decline was apparently due to low reproductive output and poor recruitment, although the reasons for these are unknown (AGFD 1999g). Possible threats to this species include degradation of habitat due to livestock grazing, off-road vehicle activity, recreation (camping and hiking), Border Patrol activities, utility corridor and road construction/maintenance, and home building (AGFD 1999g).

Because of the recent decline in monitored populations and drought conditions noted in 2002, additional surveys will be conducted prior to construction in potential supine bean habitat. If populations of this species are found in the vicinity of construction, consultation with USFS biologists will be initiated to minimize impacts. Development of the proposed TEP transmission line is likely to have an impact on this species. However, once additional surveys are completed, impacts are likely to be limited to individual plants and not whole populations. Therefore, impacts are not likely to result in a trend toward federal listing or loss of viability.

Sweet acacia (*Acacia smallii*)

The sweet acacia is a woody perennial spiny shrub or small tree found in Texas, Arizona, and California south to Argentina. Within Arizona, this species is found in the Baboquivari Mountains of Pima County and Sycamore Canyon and Atascosa Mountains of Santa Cruz County. Sweet acacia are typically found in the lower slopes of canyons of riparian areas in desert grassland communities from elevations ranging from 1,067 to 1,219 m (3,500 – 4,000 ft) (AGFD 1992).

Population trends for the sweet acacia are unknown (AGFD 1992). The proposed TEP transmission line may cross potential sweet acacia habitat; however, construction within riparian habitats will be minimized to the greatest extent possible. Placement of the transmission line may impact individual plants, however because of the linear nature of the project, only a small percentage of the population within the project area may be impacted. Furthermore, populations of this species occur outside of the project area. Therefore, impacts are not likely to result in a trend toward federal listing or loss of viability.

Thurber hoary pea (*Tephrosia thurberi*)

The Thurber hoary pea is a perennial shrub that occurs in southern Arizona and Mexico (northern Sonora and southwestern Chihuahua). Within Arizona, this plant can be found in Cochise, Santa Cruz, and Pima counties. On the Nogales RD, Thurber hoary pea plants are found in the Santa Rita and Atascosa mountains. This species typically occurs

on rocky slopes among oaks, pines, junipers, manzanitas, open hilltops, and grasslands at elevations between 1,067 and 2,134 m (3,500 – 7,000 ft) (AGFD 1999h).

There is no information on the potential effects of land use activities, such as utility placement, on the population status of Thurber hoary pea (AGFD 1999h). Placement of the transmission line may impact individual plants, however because of the linear nature of the project, only a small percentage of the population within the project area may be impacted. Furthermore, populations of this species occur in isolated mountain ranges throughout southern Arizona. Therefore, impacts to this species are not likely to result in a trend toward federal listing or loss of viability.

Thurber's morning-glory (*Ipomoea thurberi*)

Thurber's morning-glory are perennial herbaceous vines that are found in southern Arizona and Mexico (Chihuahua and Sonora). Within Arizona, this plant is found in the Huachuca and Mule Mountains of Cochise County, the Santa Rita Mountains of Pima County, and in the vicinity of Nogales, the Canelo Hills, and the Patagonia and Atascosa/Pajarito mountains of Santa Cruz County. Habitat in Arizona typically consists of rocky hillsides and canyon slopes in madrean evergreen woodland and semi-desert grassland communities in elevations between 1,158 and 1,570 m (3,800 – 5,150 ft) (AGFD 2000i). On the Nogales RD, this morning glory has been found in the vicinity of Peña Blanca Lake, east of Peñasco Canyon, and Bear Valley (T. Newman, CNF, pers. comm., 20 August 2002).

There is no information on the potential effects of land use activities, such as utility placement, on the population status of Thurber's morning-glory (AGFD 2000i). Placement of the transmission line may impact individual plants, however because of the linear nature of the project, only a small percentage of the population within the project area may be impacted. Furthermore, populations of this species occur in isolated mountain ranges throughout southern Arizona. Therefore, impacts to this species are not likely to result in a trend toward federal listing or loss of viability.

Virlet paspalum (*Paspalum virletti*)

The virlet paspalum is a perennial grass found in southeastern Arizona and Mexico (Sonora and San Luis Potosi). Within Arizona, this grass is found in the Huachuca Mountains of Cochise County, and in the Pajarito Mountains and Sycamore Canyon of Santa Cruz County. This grass is found in sandy soils of canyon bottoms in semi-desert grassland communities and grassy areas within madrean evergreen woodland communities at elevations ranging from 1,067 to 1,737 m (3,500 – 5,700 ft) (AGFD 1999i). In the Nogales RD, the only known location for this grass is in Sycamore Canyon growing in a sandy canyon bottom (T. Newman, CNF, pers. comm., 20 August 2002).

This species is rare in Arizona, where it is known from only 2 widely separated populations. There is no information on the potential effects of land use activities, such as utility placement, on the population status of virlet paspalum (AGFD 1999i). Known locations of this plant occur outside of the proposed TEP transmission line corridor; therefore, placement of the line is not likely to impact the virlet paspalum.

Weeping muhly (Sycamore Canyon muhly) (*Muhlenbergia xerophila*)

Weeping muhly is a perennial herbaceous grass found only in southern Arizona. Populations occur in the Santa Catalina, Rincon, Santa Rita, Tumacacori, and Baboquivari mountains of Pima County, and in Sycamore Canyon within the Pajarito Mountains of Santa Cruz County. Weeping muhly most often grow in crevices of cliffs, bedrock, and other rocks along canyon bottoms. This grass is also known from rocky canyon slopes in oak, pine-oak, and riparian woodlands at elevations between 1,073 and 1,829 m (3,520 – 6,000 ft) (AGFD 1999j).

There is no information on the potential effects of land use activities, such as utility placement, on the population status of weeping muhly (AGFD 1999j). Placement of the transmission line may impact individual plants, however because of the linear nature of the project, only a small percentage of the population within the project area may be impacted. Furthermore, populations of this species occur in isolated mountain ranges throughout southern Arizona. Therefore, impacts to this species are not likely to result in a trend toward federal listing or loss of viability.

Wiggins milkweed vine (*Metastelma mexicanum*)

Wiggins milkweed vine is a perennial herbaceous vine with a woody base found in southeastern Arizona to southern Sonora, Mexico. Within Arizona, this vine occurs around the Nogales and Ruby areas, Sycamore Canyon area, and Patagonia Mountains of Santa Cruz County, and Baboquivari, Coyote, and Catalina mountains of Pima County. This vine is typically found on open slopes within open oak woodland on granite soils of juniper flats at elevations between 1,067 and 1,554 m (3,500 – 5,100 ft) (AGFD 2000j). Wiggins milkweed vine has been found in several locations within the Nogales RD (T. Newman, CNF, pers. comm., 20 August 2002).

Populations of Wiggins milkweed vine within Arizona appear to be stable. This vine depends on surrounding vegetation for microhabitat and will be affected by any disturbance to area habitat (AGFD 2000j). Placement of the transmission line may impact individual plants, however because of the linear nature of the project, only a small percentage of the population within the project area may be impacted. Furthermore, populations of this species occur in isolated mountain ranges throughout southern Arizona. Therefore, impacts to this species are not likely to result in a trend toward federal listing or loss of viability.

Woolly fleabane (*Laennecia eriophylla*)

Woolly fleabane is a perennial herb found in southeastern Arizona and northern Mexico (Sonora and Chihuahua). In Arizona, woolly fleabane occurs in the Atascosa Mountains, Pajarito Mountains, Santa Rita Mountains, Canelo Hills, and in the vicinity of Sonoita Creek in Santa Cruz County. This species is typically found in gravelly soil of rocky slopes and ridges with dense grass cover in semi-desert grassland, dry oak woodland, and pine-oak woodland communities at elevations between 1,292 and 1,722 m (4,240 – 5,650

ft) (AGFD 1999k). There are known locations of wooly fleabane in the Nogales RD (T. Newman, CNF, pers. comm., 20 August 2002).

Population sizes of this plant are usually very small, with typically no more than 40 plants found in any of the populations known from Arizona. Population numbers fluctuate with the amount and timing of summer rains from year to year. This species was probably more common before its habitat was altered by excessive grazing (AGFD 1999k). Known locations of this plant and potential habitat occur outside of the proposed TEP transmission line corridor; therefore, placement of the line is not likely to impact the wooly fleabane.

3.2 INVERTEBRATES

Arizona metalmark (*Calephelis rawsoni arizonensis*)

The Arizona metalmark is a small, brown butterfly with bands of blue metallic markings on the upper and underside of the body. This butterfly occurs in Arizona, and from the Animas Mountains in southwestern New Mexico southward to Sonora, Mexico. The southern limits of its range are poorly defined to date. In Arizona, this species is known from as far north as Gila County then southward through Graham, Cochise, Pima, and Santa Cruz counties in most of the mountains therein. Arizona metalmark butterflies occur mostly above the desert floor in mountain foothills. Within these mountains, it is found in riparian canyons in oak woodland or more arid regions at elevations from 716 to 1,676 m (2,350 – 5,500 ft). Canyons with standing water for a major portion of the year appear to contain populations of this species as long as *Agave* spp. are present for larvae development (AGFD 2001a). There is no information on the potential effects of land use activities, such as utility placement, on the population status of Arizona metalmark (AGFD 2001a).

Placement of the transmission line may indirectly impact individuals of this species through habitat modification, however because the species is widely distributed across southern Arizona, only a small percentage of Arizona metalmarks may be impacted. Furthermore, transplanting of agave plants also will minimize impacts. Impacts are not likely to result in a trend toward federal listing or loss of viability.

3.3 BIRDS

American peregrine falcon (*Falco peregrinus anatum*)

The American peregrine falcon subspecies is a medium-sized raptor that nests from central Alaska south to Baja California, Sonora, and the highlands of Central Mexico. Within Arizona, this raptor breeds wherever sufficient prey is available near cliffs. These raptors are rare or absent as breeders in the southwestern quarter of Arizona. Optimum habitat for peregrine falcons consists of steep, sheer cliffs overlooking woodlands, riparian areas, or other habitats supporting avian prey species in abundance. These raptors may also be found in less optimal habitat consisting of small broken cliffs in ponderosa pine forests or large sheer cliffs in very xeric areas. The presence of an open expanse is critical. American peregrine falcons can be found at elevations ranging from

122 to 2,743 m (400 – 9,000 ft) (Glinski 1998, AGFD 1998f). Peregrine falcon nests were found on Ramanote Peak and along Sycamore Canyon (CNF 2000). Both these nests are at least 1.6 km (1 mi) from the proposed ROW. In 2002, another nest was found on Castle Rock, which is within the MSO PAC and within 0.3 km (0.18 mi) of proposed structures. The seasonal restrictions in effect for MSO (SECTION 1.4) will prevent breeding season disturbance of peregrines on Castle Rock.

American peregrine falcons have been found in great numbers in Arizona as well as in areas that will have formerly been considered marginal habitat. This trend suggests that populations in Arizona may have reached levels saturating the optimal habitat available (AGFD 1998f). Placement of the proposed transmission line is not near known nesting sites for peregrine falcons. If new nest sites are encountered during construction, conservation measures will be developed in coordination with CNF biologists to prevent adverse effects. Therefore, placement of the transmission line will not impact this species.

Five-striped sparrow (*Aimophila quinquestriata*)

The five-striped sparrow is found in western portions of northern Sinaloa and Sonora, Mexico and the southeastern most portions of Arizona. This sparrow is primarily found in Mexico, but its range reaches into southeastern Arizona. Here, it is rarely found during breeding season, and there are only a few winter records. Five-striped sparrow habitat is highly specialized, consisting of tall, dense shrubs on rocky, semi-desert hillsides and canyon slopes (New Mexico Game and Fish Department and the Fish and Wildlife Information Exchange 2000). Within the Nogales RD, this sparrow has been recorded within Sycamore Canyon (T. Newman, CNF, pers. comm., 20 August 2002).

Populations of five-striped sparrow have declined because of habitat loss, fragmentation, and degradation (New Mexico Game and Fish Department and the Fish and Wildlife Information Exchange 2000). The proposed TEP transmission line will not cross Sycamore Canyon where these sparrows have been observed. This species is not likely to be affected by the proposed placement of a transmission line within the Nogales RD.

Northern gray hawk (*Asturina nitida maxima*)

The gray hawk is a medium-sized raptor with a gray back, black tail with 2 or 3 white bands, and a finely barred gray and white chest, abdomen, and thighs (Glinski 1998). The gray hawk prefers Sonoran riparian deciduous forest and woodland plant communities and can be found along the Santa Cruz and San Pedro rivers, Sonoita Creek, and Sopori Wash. This species also has been reported from the Hassayampa and Salt rivers. This hawk species is migratory and usually arrives in Arizona in mid-March and returns south during winter months (AGFD 2000k). Gray hawks prefer cottonwood, mesquite, and hackberry woodlands with a prey base of lizards, especially the whiptail lizard (*Cnemidophorus* spp.).

The current population trend for gray hawks is considered stable by the AGFD (2000k). Potential nesting habitat exists near the proposed TEP transmission line corridor in Peck

Canyon. Individual gray hawks may be impacted by noise from construction activity related to transmission line placement. However, because of the distance of the proposed action from suitable habitat in Peck Canyon, any increase in noise will be marginal. Furthermore, only a small percentage of the population may be impacted. Therefore, impacts are not likely to result in a trend toward federal listing or loss of viability.

Western yellow-billed cuckoo (*Coccyzus americanus occidentalis*)

The western yellow-billed cuckoo is a long and slender bird with short, dark legs that nests from southern California through the northeastern United States, south through the United States to the Florida Keys, Central America and southern Baja California, Mexico. This species winters from South America to central Argentina and Uruguay. Within Arizona, western yellow-billed cuckoo are found in southern and central Arizona and the extreme northeast portion of the state. This species is typically found in streamside areas with cottonwood, willow groves, and larger mesquite bosques (AGFD 1998g). This species has been observed in Sopori Wash and Sycamore, Peck, and Peña Blanca canyons (AGFD 1998g; CNF 2000; P. Titus, T. Furgason, SWCA, pers. comm. 16 October 2002).

Populations of western yellow-billed cuckoo have been reduced; a general decline is occurring in all areas with known populations (AGFD 1998g). This species is sensitive to habitat fragmentation and degradation of riparian woodlands due to agricultural and residential development (Hughes 1999). The proposed transmission line may cross potential cuckoo habitat; however, construction within riparian habitats will be minimized to the greatest extent possible. Placement of the transmission line may impact individuals of this species, however because of the linear nature of the project, only a small percentage of the population within the project area may be impacted. Furthermore, populations of this species occur outside of the project area. Therefore, impacts are not likely to result in a trend toward federal listing or loss of viability.

3.4 REPTILES AND AMPHIBIANS

Giant spotted whiptail (*Cnemidophorus burti strictogrammus*)

The giant spotted whiptail is a long, slender lizard found in southeastern Arizona, extreme southwest New Mexico, and northern Sonora, Mexico. Within southeastern Arizona, this lizard is found in Cochise County; the Santa Catalina, Santa Rita, Baboquivari, and Pajarito mountains and in the vicinity of Oracle in Pima County; and in Pinal County. Giant spotted whiptail lizards inhabit mountain canyons, arroyos, and mesas in arid and semi-arid regions, entering lowland deserts along stream courses. They are found in dense shrubby vegetation, often among rocks near permanent and intermittent streams at elevations ranging from near sea level to 1,372 m (4,500 ft). Open areas of bunch grass within these riparian habitats are also occupied (AGFD 2001b).

Giant spotted whiptail populations are thought to be stable and some populations are locally abundant even though this species is limited in distribution (AGFD 2001b). Because the known populations occur outside the project area, the proposed transmission line will have no significant effect on the population status of the giant spotted whiptail.

Lowland leopard frog (*Rana yavapaiensis*)

The lowland leopard frog is found in low elevations in the drainage of the lower Colorado River and its tributaries in Nevada, California, Arizona, New Mexico, northern Sonora and extreme northeast Baja California, Mexico (probably extirpated from California and Nevada). Within Arizona, this frog has been found in the Virginia River drainage in the extreme northwestern part of the state, in the Colorado River near Yuma, and west, central, and southeast Arizona south of the Mogollon Rim. This frog frequents desert, grassland, oak, and oak-pine woodland in permanent pools of foothill streams, rivers, and permanent stock tanks. They typically stay close to water at elevations ranging from 244 to 1,676 m (800 – 5,500 ft) (AGFD 1997b). Within the Nogales RD, this frog has been recorded in Pesquiera and Alamo canyons, California Gulch, Adobe, Temporal Gulch, Big Casa Blanca, Box Canyon, and Gardner Canyon (T. Newman, CNF, pers. comm., 20 August 2002).

Lowland leopard frog populations are considered stable in central Arizona but declining in southeast Arizona, and populations have been extirpated from southwestern Arizona. Potential threats to this species are manipulation to major watercourses, water pollution, introduced species (fish, bullfrogs, and crayfish), heavy grazing, and habitat fragmentation (AGFD 1997b). Because no construction will occur within perennial aquatic habitats and known populations occur outside project area, the proposed transmission line will have no significant effect on the population status of the lowland leopard frog.

Mexican garter snake (*Thamnophis eques megalops*)

The Mexican garter snake ranges from southeastern Arizona and extreme southwestern New Mexico, southward into the highlands of western and southern Mexico, to Oaxaca. Within Arizona, this snake occurs in the southeast corner of the state from the Santa Cruz Valley east and generally south of the Gila River. Valid records (post 1980) have recorded this snake in the San Rafael and Sonoita grasslands area and from Arivaca. Mexican garter snakes are most abundant in densely vegetated desert grassland habitat surrounding cienegas, cienega-streams, stock tanks, and in or near water along streams in valley floors and generally open areas, but not in steep mountain canyon stream habitat. This snake is generally found at elevations ranging from 914 to 1,524 m (3,000 – 5,000 ft) but may reach elevations of 2,591 m (8,500 ft) (AGFD 2001c).

Populations of Mexican garter snakes are decreasing, with extirpations at several localities since 1950 as habitat has changed and introduced predators have invaded. Management concerns for this species include predation by introduced bullfrogs and predatory fishes, urbanization and lowered water tables, and habitat destruction, including that due to overgrazing (AGFD 2001c). Because no construction will occur within perennial aquatic habitats and construction within riparian habitats will be minimized, the proposed transmission line will have no significant effect on the population status of the Mexican garter snake.

Western barking frog (*Eleutherodactylus augusti cactorum*)

The western barking frog is a secretive terrestrial frog found in extreme southern Arizona, southeast New Mexico, and central Texas south to the Isthmus of Tehuantepec. In Arizona, this frog historically occurred in Pima and Santa Cruz counties within the Santa Rita and Pajarito mountains. Habitat consists of rocky hillsides of canyons in woodland vegetation at elevations between 1,158 and 2,134 m (3,800 – 7,000 ft). Permanent water is not a necessary component of western barking frog habitat. There are very few records of this species in Arizona, and none have been recorded within the Nogales RD (AGFD 1995b).

There is no information on the potential effects of land use activities, such as utility placement, on the population status of western barking frogs (AGFD 1995b). Because known populations occur outside the project area, the proposed transmission line will have no significant effect on the population status of the western barking frog and is not likely to result in a trend toward listing or loss of viability.

3.5 MAMMALS

Cave myotis (*Myotis velifer*)

The cave myotis is a large bat found in the southwestern half of Arizona and the immediate adjacent parts of California, Nevada, New Mexico, and the northern third of Sonora, Mexico. Within Arizona, this bat is found south of the Mogollon Plateau from Lake Mohave, Burro Creek, Montezuma Well, San Carlos Apache Reservation, and the Chiricahua Mountains south to Mexico. Cave myotis have not been recorded in the extreme southwestern part of the state and are found in small numbers in southeastern Arizona in the winter. This bat typically prefers desertscrub habitats of creosote, brittlebush, paloverde, and cacti but they sometimes can be found up in pine-oak communities. Cave myotis roost in caves, tunnels, mineshafts, under bridges, and sometimes buildings within a few kilometers of a water source (AGFD 1997c).

Cave myotis colonies are vulnerable at the roost sites, especially maternity roosts, because they congregate in large numbers (AGFD 1997c). The proposed TEP transmission line will not cross near known roost sites. Potential foraging habitat may be disturbed during development of the transmission line; however, these disturbances will be isolated and widely distributed. Furthermore, populations of this species occur throughout southern Arizona. Therefore, impacts will not likely result in a trend toward federal listing or loss of viability of the cave myotis.

Southern pocket gopher (*Thomomys umbrinus intermedius*)

The southern pocket gopher is a small gopher found in extreme southeastern Arizona and southwestern New Mexico, south into Mexico. Within Arizona, this gopher is found primarily in the southern most portion of the state in the oak belt of the Santa Rita, Patagonia, Atascosa, Pajarito, and Huachuca mountains. Southern pocket gophers have been found at Peña Blanca Spring in gravelly soil along a broad wash. Elsewhere, this species is generally found on rocky slopes within open oak woodlands in the lower parts of mountain ranges from 1,372 to 2,743 m (4,500 – 9,000 ft) in elevation. There has been

only 1 record for the southern pocket gopher within the Nogales RD, specifically at Peña Blanca Canyon in the Atascosa/Pajarito mountains. However, it is suspected that this species has a much wider range (AGFD 1998h).

There is no information on the potential effects of land use activities, such as utility placement, on the population status of southern pocket gopher (AGFD 1998h). Placement of the transmission line may impact individuals of this species, however because of the linear nature of the project, only a small percentage of the population within the project area may be impacted. Furthermore, populations of this species occur outside of the project area. Therefore, impacts are not likely to result in a trend toward federal listing or loss of viability.

4.0 BLM SENSITIVE SPECIES

Criteria for BLM Sensitive species include those that are:

1. Under status review by the USFWS, or
2. Whose numbers are declining so rapidly that Federal listing may become necessary, or
3. With typically small and widely dispersed populations,
4. Those inhabiting ecological refugia or other specialized or unique habitats.

The potential impacts to BLM Sensitive species were determined based on the habitat conditions within the BLM lands crossed by the proposed action, the life history of the species, and the proposed construction methods. Only those species that have a potential of occurring on or near the BLM parcel were evaluated. The 13 BLM Sensitive species evaluated were identified in the BLM Sensitive species list for Arizona (Instruction Memorandum No. AZ-2000-018) dated 21 April 2000 and are listed in Table 4.

TABLE 4. SUMMARY OF EFFECTS ON BUREAU OF LAND MANAGEMENT SENSITIVE SPECIES.		
COMMON NAME <i>Scientific Name</i>	EFFECTS DETERMINATION	JUSTIFICATION
Balloonvine <i>Cardiospermum corindum</i>	May impact individuals of this species, but is not likely to result in a trend toward federal listing or loss of viability.	<ul style="list-style-type: none"> Only small percentage of total population within project area may be impacted. Other viable populations occur outside of project area.
False grama <i>Cathastecum erectum brevifolium</i>	May impact individuals of this species, but is not likely to result in a trend toward federal listing or loss of viability.	<ul style="list-style-type: none"> Only small percentage of total population within project area may be impacted. Other viable populations occur outside of project area.
Tumamoc globeberry <i>Tumamoca macdougallii</i>	May impact individuals of this species, but is not likely to result in a trend toward federal listing or loss of viability.	<ul style="list-style-type: none"> Minimal impacts to riparian habitat. Only small percentage of total population within project area may be impacted. Other viable populations occur outside of project area.
Loggerhead shrike <i>Lanius ludovicianus</i>	May impact individuals of this species, but is not likely to result in a trend toward federal listing or loss of viability.	<ul style="list-style-type: none"> Only small percentage of total population within project area may be impacted. Populations of this species occur throughout southern Arizona.
Rufous-winged sparrow <i>Aimophila carpalis</i>	May impact individuals of this species, but is not likely to result in a trend toward federal listing or loss of viability.	<ul style="list-style-type: none"> Only small percentage of total population within project area may be impacted. Other viable populations occur outside of project area.

TABLE 4 (CONTINUED). SUMMARY OF EFFECTS ON BUREAU OF LAND MANAGEMENT SENSITIVE SPECIES.		
COMMON NAME <i>Scientific Name</i>	EFFECTS DETERMINATION	JUSTIFICATION
Western burrowing owl <i>Athene curvicularia hypugea</i>	May impact individuals of this species, but is not likely to result in a trend toward federal listing or loss of viability.	<ul style="list-style-type: none"> Only small percentage of total population within project area may be impacted. Populations of this species occur throughout southwestern U.S.
Texas horned lizard <i>Phrynosoma cornutum</i>	No Impacts.	<ul style="list-style-type: none"> Known populations occur outside project area.
Big free-tailed bat <i>Nyctinomops macrotis</i>	May impact individuals of this species, but is not likely to result in a trend toward federal listing or loss of viability.	<ul style="list-style-type: none"> No known roosts within project area. Only small percentage of foraging habitat within project area may be impacted. Populations of this species occur throughout southern Arizona.
California leaf-nosed bat <i>Macrotus californicus</i>	May impact individuals of this species, but is not likely to result in a trend toward federal listing or loss of viability.	<ul style="list-style-type: none"> No known roosts within project area. Only small percentage of foraging habitat within project area may be impacted. Populations of this species occur throughout southern Arizona.
Fringed myotis <i>Myotis thysandodes</i>	May impact individuals of this species, but is not likely to result in a trend toward federal listing or loss of viability.	<ul style="list-style-type: none"> No known roosts within project area. Only small percentage of foraging habitat within project area may be impacted. Populations of this species occur throughout southern Arizona.
Pocketed free-tailed bat <i>Nyctinomops femorosaccus</i>	May impact individuals of this species, but is not likely to result in a trend toward federal listing or loss of viability.	<ul style="list-style-type: none"> No known roosts within project area. Only small percentage of foraging habitat within project area may be impacted. Populations of this species occur throughout southern Arizona.
Spotted bat <i>Euderma maculatum</i>	May impact individuals of this species, but is not likely to result in a trend toward federal listing or loss of viability.	<ul style="list-style-type: none"> No known roosts within project area. Only small percentage of foraging habitat within project area may be impacted. Populations of this species occur throughout southern Arizona.
Underwood's mastiff bat <i>Eumops underwoodi</i>	May impact individuals of this species, but is not likely to result in a trend toward federal listing or loss of viability.	<ul style="list-style-type: none"> No known roosts within project area. Only small percentage of foraging habitat within project area may be impacted. Populations of this species occur throughout southern Arizona.

4.1 PLANTS

Balloonvine (*Cardiospermum corindum*)

This perennial vine is widely distributed in tropical and subtropical regions and is known from the Coyote Mountains in Pima County (Kearny and Peebles 1960). Because potential habitat for this species is widespread, placement of the transmission line may impact individual plants. However because of the linear nature of the project, only a

small percentage of the population within the project area may be impacted. Furthermore, populations of this species occur outside of the project area. Therefore, impacts are not likely to result in a trend toward federal listing or loss of viability.

False grama (*Cathestecum erectum (brevifolium)*)

False grama is a perennial, drought-tolerant grass found on dry hills and plains of Southern Arizona and Northern Mexico. Placement of the transmission line may impact individual plants, however because of the linear nature of the project, only a small percentage of the population within the project area may be impacted. Furthermore, populations of this species occur outside of the project area. Therefore, impacts to this species are not likely to result in a trend toward federal listing or loss of viability.

Tumamoc globeberry (*Tumamoca macdougalii*)

This perennial vine occurs in shade of nurse plants along sandy washes below ~914 m (3,000 ft) in elevation. The proposed transmission line may cross potential habitat for this species; however, construction within riparian habitats will be minimized to the greatest extent possible. Placement of the transmission line may impact individual plants, however because of the linear nature of the project, only a small percentage of the population within the project area may be impacted. Furthermore, populations of this species occur outside the project area. Therefore, impacts to this species are not likely to result in a trend toward federal listing or loss of viability.

4.2 BIRDS

Loggerhead shrike (*Lanius ludovicianus*)

The loggerhead shrike occurs in open country with scattered trees and shrubs, savanna, desertscrub and occasionally open woodland (AGFD 2002). In Arizona, this species usually summers throughout open parts of the state below the Transition Zone and is also periodically found along the Mexican border west of Baboquivari Mountains (Phillips et al. 1983). Because habitat for this species is widely distributed, placement of the transmission line may impact this species. However because of the linear nature of the project, only a small percentage of the population within the project area may be impacted. Furthermore, populations of this species occur throughout southern Arizona. Therefore, impacts are not likely to result in a trend toward federal listing or loss of viability.

Rufous-winged sparrow (*Aimophila carpalis*)

The rufous-winged sparrow is classified as a migratory bird and is a resident of eastern Pima County, including Avra Valley, and was once thought to be extirpated in Arizona due to overgrazing but was rediscovered in the Tucson Area in 1936. Rufous-winged sparrows generally use habitats characterized by scattered low shrubs and trees, which provide cover and foraging areas during mid-summer days. Many of these areas contain significant grassland components. Threats to the species include urban development, overgrazing, and exotic species, all of which result in losses of grassland communities utilized by this species (Pima County 2001). Because habitat for this species is widely distributed, placement of the transmission line may impact this species. However

because of the linear nature of the project, only a small percentage of the population within the project area may be impacted. Furthermore, populations of this species occur outside the project area. Therefore, impacts are not likely to result in a trend toward federal listing or loss of viability.

Western burrowing owl (*Athene cunicularia hypugea*)

The Western burrowing owl inhabits heavily grazed tracts of mixed-grass prairie, particularly where there are burrows created by large rodents, such as prairie dogs and Richardson ground squirrels. Distribution extends from southern Canada through the western United States to South America. Arizona is 1 of 3 states that provide important wintering areas for this species (USGS 2003). Because habitat for this species is widely distributed, placement of the transmission line may impact this species. However because of the linear nature of the project, only a small percentage of the population within the project area may be impacted. Furthermore, populations of this species occur throughout the southwestern United States. Therefore, impacts are not likely to result in a trend toward federal listing or loss of viability.

4.3 REPTILES AND AMPHIBIANS

Texas horned lizard (*Phrynosoma cornutum*)

The Texas horned lizard occurs from Kansas to extreme southeastern Arizona and lives mainly in sandy areas of deserts, grasslands, prairies, and scrublands (Bartlett and Bartlett 1999) where it often inhabits abandoned animal burrows (Bockstanz 1998). Because known populations occur outside of the project area, the proposed transmission line will have no significant effect on the population status of this species.

4.4 MAMMALS

Big free-tailed bat (*Nyctinomops macrotis*)

Distribution of the big free-tailed bat occurs from the southwestern United States southward through the Caribbean, Central America, and into the northern part of South America. Northern populations are known to migrate to southern Arizona and Mexico in the fall, yet this species is widely scattered throughout Arizona during the spring and summer too. In Arizona, this bat has been found in pinyon-juniper, Douglas-fir, and Sonoran desertscrub habitats, but it is believed that these locations are foraging sites. Preferred roosting sites include rock crevices and fissures of mountain cliffs in rugged, rocky areas of desertscrub habitat (AGFD 1993, Harvey et al. 1999). The proposed TEP transmission line will not cross near known roost sites, but potential foraging habitat may be disturbed during development of the transmission line; however, these disturbances will be isolated and widely distributed. Furthermore, populations of this species occur throughout southern Arizona. Therefore, impacts will not likely result in a trend toward federal listing or loss of viability of the big free-tailed bat.

California leaf-nosed bat (*Macrotus californicus*)

Distribution of the California leaf-nosed bat in the United States spans southern California, southern Nevada, and southwestern Arizona and extends southward into

Mexico, to the southern tip of Baja California, northern Sinaloa, and southwestern Chihuahua. This bat lives predominantly in Sonoran and Mohave desertscrub habitats, but is occasionally found in the Chihuahuan and Great Basin deserts. Daytime roosting sites are usually mines and caves, and nighttime roosts include open buildings, cellars, bridges, porches, and mines. These bats do not hibernate or migrate; therefore, they tend to live in the same area year after year and remain active year-round (AGFD 1993, 2001d; Harvey et al. 1999). The proposed TEP transmission line will not cross near known roost sites, but potential foraging habitat may be disturbed during development of the transmission line; however, these disturbances will be isolated and widely distributed. Furthermore, populations of this species occur throughout southern Arizona. Therefore, impacts will not likely result in a trend toward federal listing or loss of viability of the California leaf-nosed bat.

Fringed myotis (*Myotis thysanodes*)

Distribution of the fringed myotis ranges from southern British Columbia, Canada southward throughout the western United States, and down to southern Mexico. It occurs in a variety of habitats – from desertscrub to oak and pinyon woodlands to spruce-fir forests. Roosting sites include caves, mines, and buildings. These bats tend to roost in tight clusters and may change locations periodically in response to thermoregulatory needs (AGFD 1993, Harvey et al. 1999). The proposed TEP transmission line will not cross near known roost sites, but potential foraging habitat may be disturbed during development of the transmission line; however, these disturbances will be isolated and widely distributed. Furthermore, populations of this species occur throughout southern Arizona. Therefore, impacts will not likely result in a trend toward federal listing or loss of viability of the fringed myotis.

Pocketed free-tailed bat (*Nyctinomops femorosaccus*)

The pocketed free-tailed bat ranges from the southwestern United States (including southern California, Arizona, and New Mexico, and the Trans-Pecos region of Texas), south into Mexico through Baja, Sonora, Durango, and Jalisco to, at least, Michoacan. This bat can be found in the arid lowlands of the desert Southwest, where it roosts in crevices and caves of rugged cliffs, slopes, and rock outcrops (AGFD 1993, Harvey et al. 1999). The proposed TEP transmission line will not cross near known roost sites. Potential foraging habitat may be disturbed during development of the transmission line; however, these disturbances will be isolated and widely distributed and will not likely result in a trend toward federal listing or loss of viability of this species.

Spotted bat (*Euderma maculatum*)

Distribution of the spotted bat ranges throughout centralwestern North America, from southcentral British Columbia down to southern Mexico. In Arizona, its habitat ranges from low desert areas in the Southwest to high desert and riparian habitats in the northwestern part of the state. This bat has also been documented in conifer forests in northern Arizona. Roosting sites are often situated in rock crevices on high cliffs (AGFD 1993, Harvey et al. 1999). The proposed TEP transmission line will not cross near known roost sites, but potential foraging habitat may be disturbed during development of the transmission line; however, these disturbances will be isolated and widely distributed.

Furthermore, populations of this species occur throughout southern Arizona. Therefore, impacts will not likely result in a trend toward federal listing or loss of viability of the spotted bat.

Underwood's mastiff bat (*Eumops underwoodi*)

The range of Underwood's mastiff bat is limited, from south-central Arizona, into the arid lowlands of Sonoran and western Mexico, and into Honduras. It is believed to be a year-round resident of Arizona, ranging from the Baboquívari Mountains down to Organpipe National Monument. This bat prefers Sonoran desertscrub and mesquite/grassland plant communities. Roosting tends to occur in crevices along steep cliffs and sometimes in the cracks of buildings (AGFD 1993). The proposed TEP transmission line will not cross near known roost sites, but potential foraging habitat may be disturbed during development of the transmission line; however, these disturbances will be isolated and widely distributed. Furthermore, populations of this species occur throughout southern Arizona. Therefore, impacts will not likely result in a trend toward federal listing or loss of viability of this species.

5.0 AGFD WILDLIFE OF SPECIAL CONCERN

AGFD was consulted in regards to state listed special status species and habitats that may be affected by the proposed action. Several state listed special status species and overall wildlife habitat may be affected by the proposed action. The AGFD mission is to conserve, enhance, and restore Arizona's diverse wildlife resources and habitats through aggressive protection and management programs. Continued consultation and input from AGFD will ensure that impacts of the proposed action are minimized and mitigation efforts are successful.

Listed in Table 5 are state special status species that may be found in the vicinity of the proposed action, based on AGFD's Heritage Data Management System (HDMS) (1 July 2002). Effects of the proposed action on the majority of these species will be avoided or minimized through mitigation efforts stipulated for federally listed species. However, additional mitigation is recommend for the Sonoran Desert tortoise as 5 individuals were located near the Tinaja Hills area during field surveys of the proposed ROW (HEG 2002, unpublished data).

TABLE 5. SUMMARY OF EFFECTS ON WILDLIFE OF SPECIAL CONCERN IN ARIZONA.		
COMMON NAME <i>Scientific Name</i>	EFFECTS DETERMINATION	JUSTIFICATION
Black-bellied whistling duck <i>Dendrocyna autumnalis</i>	No Impacts.	<ul style="list-style-type: none"> No construction in perennial aquatic habitats.
Crested caracara <i>Caracara cheriway</i>	No Impacts.	<ul style="list-style-type: none"> Known populations occur outside project area.
Desert tortoise - Sonoran population <i>Gopherus agassizii</i>	May impact individuals of this species, but is not likely to result in a trend toward federal listing or loss of viability.	<ul style="list-style-type: none"> Only small percentage of total potential habitat within project area may be impacted. Pre-construction surveys will minimize impacts to species.
Elegant trogon <i>Trogon elegans</i>	May impact individuals of this species, but is not likely to result in a trend toward federal listing or loss of viability.	<ul style="list-style-type: none"> Minimal impacts to riparian habitat. Only small percentage of total population within project area may be impacted. Populations of this species occur in isolated mountain ranges throughout southern Arizona.
Mexican long-tongued bat <i>Choeronycteris mexicana</i>	May impact individuals of this species, but is not likely to result in a trend toward federal listing or loss of viability.	<ul style="list-style-type: none"> Only small percentage of total potential habitat within project area may be impacted. Mitigation plantings of agaves will reduce impacts.
Great Plains narrow-mouthed toad <i>Gastrophryne olivacea</i>	May impact individuals of this species, but is not likely to result in a trend toward federal listing or loss of viability.	<ul style="list-style-type: none"> Minimal impacts to riparian habitat. Only small percentage of total population within project area may be impacted. Other viable populations occur outside of project area.

TABLE 5 (CONTINUED). SUMMARY OF EFFECTS ON WILDLIFE OF SPECIAL CONCERN IN ARIZONA.

COMMON NAME <i>Scientific Name</i>	EFFECTS DETERMINATION	JUSTIFICATION
Mexican vine snake <i>Oxibelis aeneus</i>	No Impacts.	<ul style="list-style-type: none"> Known occurrences are outside project area.
Osprey <i>Pandion haliaetus</i>	No Impacts	<ul style="list-style-type: none"> No construction in perennial aquatic habitats.
Rose-throated becard <i>Pachyramphus aglaiae</i>	No Impacts.	<ul style="list-style-type: none"> Known occurrences are outside project area.
Thick-billed kingbird <i>Tyrannus crassirostris</i>	No Impacts	<ul style="list-style-type: none"> No potential habitat within project area.
Tropical Kingbird <i>Tyrannus melancholicus</i>	May impact individuals of this species, but is not likely to result in a trend toward federal listing or loss of viability.	<ul style="list-style-type: none"> Minimal impacts to riparian habitat. Only small percentage of total population within project area may be impacted. Other viable populations occur outside of project area.

Black-bellied whistling duck (*Dendrocyna autumnalis*)

The black-bellied whistling duck is "goose-like" with a long neck and long pink legs. This species has a cinnamon or chestnut breast and back with a black belly and bright coral-red bill. The total range for this species is from the Gulf coast and lower Rio Grande Valley of Texas and central Arizona south through Mexico, Central America to southern Brazil. In Arizona, the range for the black-bellied whistling duck is southeastern and central Arizona. Black-bellied whistling ducks are commonly seen in the Santa Cruz Valley, particularly in ponds near and around Nogales. The habitat for this species consists of the banks of rivers, lakes, ponds, riparian areas, and stock tanks (Brown 1985).

Because of habitat loss and apparent population declines from historic levels, the black-bellied whistling duck has been placed on the AGFD Threatened Native Wildlife of Arizona List as a candidate species. This species appears to be increasing in Arizona in urban settings at man-made ponds and at sewage treatment plants. It also appears to be stable at some private ranch ponds, which tend to be isolated from hunting pressure (Corman 1994).

Because no construction will occur in perennial aquatic habitats, the proposed transmission line will have no effect on the population status of the black-bellied whistling duck.

Crested caracara (*Caracara cheriway*)

The crested caracara is a medium sized raptor with bold black and white plumage and a bright yellow-orange face and legs. The crested caracara ranges from southern Arizona

and northern Mexico to Tierra del Fuego. In the United States, it occurs only along the southern border in Texas and Arizona, and in Florida, where there is an isolated population in the south-central peninsula. In Arizona, their range extends up from San Miguel in the Baboquivari Valley north to Quijotoa, Sells, and Coyote Pass. This raptor occurs regularly on the Tohono O'odham Indian Reservation. Small groups of crested caracara are seen in Sasabe and south of the Mexican border near Sonoyta, Sonora. This raptor is found in open habitats, typically grassland, prairie, pastures, or desert with scattered taller trees, shrubs, or cacti. The crested caracara is found in areas characterized by low-profile ground vegetation and scattered tall vegetation. Specifically in Arizona, vegetation consists of saguaro, mesquite, palo verde, cholla and acacia (Morrison 1996).

Arizona populations of crested caracara on the Tohono O'odham Reservation are likely stable because few threats exist. Reports of individual, and in some cases groups, of this raptor outside of the reservation indicate that its range within Arizona is probably as extensive as it was historically. No apparent threat currently exists to Arizona populations; however, the AGFD has listed the crested caracara as a threatened native wildlife. This species is considered vulnerable if habitat conditions worsen (Morrison 1996).

Habitat surveys did not detect the presence of any bird of prey nests along the corridor. Furthermore, no known populations of this species occur within the project area. Therefore, the proposed action will have no effect on the population status of the crested caracara.

Desert tortoise (Sonoran) (*Gopherus agassizii*)

The Sonoran Desert tortoise ranges from northern Sinaloa, Mexico to southern Nevada and southwestern Utah, and from southcentral California east to southeastern Arizona. The desert tortoise is divided into 2 populations for purposes of the Endangered Species Act. The threatened Mojave population occurs north and west of the Colorado River and the unlisted Sonoran population occurs south and east of the Colorado River. Within Arizona, the Sonoran Desert tortoise is found south and east of the Colorado River from Mojave County to the south, beyond the International Boundary and many scattered locations in between. The Sonoran population of the desert tortoise occurs primarily on rocky slopes and bajadas of Mojave and Sonoran desert scrub at elevations ranging from 152 to 1,615 m (500 – 5,300 ft). Burrows and shelter sites are generally below rocks and boulders, in rock crevices, under vegetation, and also in caliche caves of incised wash banks (AGFD 2001e).

Several threats to tortoise populations in the Sonoran Desert have been identified, including habitat fragmentation, habitat loss and degradation from urban and agricultural development and roads, wildfires associated with invasion of non-native grasses and forbs, illegal collection, and genetic contamination of wild populations by escaped or released captives. Although current evidence suggests that Arizona populations are stable there are substantial gaps in available data (Arizona Interagency Desert Tortoise Team 1996).

During ground surveys of the proposed transmission line corridor, 5 desert tortoise were found (HEG, unpublished data). Per recommendations of Spencer and Humphrey (1999) for any ground disturbing projects, surveys should be conducted a minimum of 48 hours prior to grading and again just prior (as it is occurring) to vegetation clearing (Desert Tortoise Council 1999). While the proposed action may have a minimal effect on the potential habitat of this species, pre-construction surveys will minimize impacts to individual tortoise and is therefore not likely to result in a trend toward listing or loss of viability.

Elegant trogon (*Trogon elegans*)

The elegant trogon is a medium sized bird with a round head, large eyes, a white band on an iridescent green breast, black face and throat, red belly and undertail coverts. The total range for this bird is from southern Arizona and New Mexico south through Mexico to southern Nicaragua to northwestern Costa Rica. In Arizona, the elegant trogon is found in sky island mountains, most commonly the Atascosa, Chiricahua, Huachuca, and Santa Rita mountains. Elegant trogons are found in riparian areas consisting of sycamore, cottonwood, and oak, and also in coniferous woodlands at elevations ranging from 1,036 to 2,073 m (3,400 – 6,800 ft) (AGFD 2001f).

Population trends for the elegant trogon are not well known. No evidence indicates population declines in any of the core canyons occupied over the past few decades. Threats to this species include degradation and loss of native riparian habitat through stream diversion, groundwater withdrawal, erosion, and overgrazing (AGFD 2001f).

Because potential habitat and known occurrences of this species are outside the project area, the proposed action will have no effect on the population status of the elegant trogon.

Great Plains narrow-mouthed toad (*Gastrophryne olivacea*)

The Great Plains narrow-mouthed toad is a small, stout toad with stubby limbs, a small pointed head with a fold of skin on the back of the head. The total range for this species is from southeastern Nebraska and Missouri south through Texas to western Mexico. Within Arizona, the Great Plains narrow-mouthed toad is found in the vicinity of Santa Cruz County, Pima County, to near Casa Grande, Arizona in Pinal County. Habitat for this species in Arizona consists of mesquite semi-desert grassland communities to oak woodland communities near riparian areas at elevations ranging from sea level to around 1,250 m (4,100 ft) (AGFD 1995c).

Population trends for the Great Plains narrow-mouthed toad are currently unknown. Populations in Arizona are at the extreme northwestern edge of the species range and distribution is limited throughout its range (AGFD 1995c). The proposed transmission line may cross potential habitat for this species; however, construction within riparian habitats will be minimized to the greatest extent possible. Placement of the transmission line may impact individuals of this species, however because of the linear nature of the

project, only a small percentage of the population within the project area may be impacted. Furthermore, populations of this species occur outside the project area. Therefore, impacts to this species are not likely to result in a trend toward federal listing or loss of viability.

Mexican long-tongued bat (*Choeronycteris mexicana*)

The Mexican long-tongued bat has a long, slender nose with a leaf-like structure on the base of the nose. The total range for this species is from southeastern Arizona, southwestern New Mexico, and California south through Central America to Venezuela. In Arizona, the Mexican long-tongued bat is found from the Chiricahua Mountains extending as far north as the Santa Catalina Mountains and west to the Baboquivari Mountains. Habitat for this bat is typically within canyons of mixed oak-conifer forests in mountains at elevations ranging from 1,082 to 2,231 m (3,550 – 7,320 ft) (AGFD 1994). This species do not congregate in sizeable maternity or bachelor colonies like *Leptonycteris* bats do (Hoffmeister 1986). They feed on nectar and pollen, especially from paniculate agaves (AGFD 1994).

Populations of Mexican long-tongued bats in Arizona appear to be highly variable (AGFD 1994) and there is no evidence of a long-term decline or any clear trend. The limitation of riparian zones and the distribution of food plants may limit populations of this species in Arizona and loss of riparian vegetation may be a greater threat to this species than human disturbance at particular roost sites (Pima County 2001). The proposed TEP transmission line will not cross near known roost sites, but potential foraging habitat may be disturbed during construction; however, these disturbances will be isolated and will impact only a small percentage of potential habitat. Furthermore, transplanting of agave plants also will minimize impacts. Impacts to this species are not likely to result in a trend toward federal listing or loss of viability.

Mexican vine snake (*Oxibelis aeneus*)

The Mexican vine snake has an elongated head, pointed snout, and is thin bodied with an ash gray to yellow-brown and tan coloring. The total range for this species is from extreme southern Arizona south to Brazil. In Arizona, this species occurs in the Tumacacori, Pajarito, and Patagonia mountains in Santa Cruz County. Habitat for the Mexican vine snake consists of brush-covered hillsides and riparian areas with sycamore, oak, walnut and wild grape trees at elevations ranging from 914 to 1,768 m (3,000 – 5,800 ft) (AGFD 1991b).

Population trends for the Mexican vine snake are currently unknown. Populations in Arizona are at the extreme northern edge of the species range and distribution is limited, with occurrences known from Sycamore Canyon (AGFD 1991b). A potential threat is the high interest by collectors for this species (AGFD 1991b). Because known occurrences of this species are outside the project area, the proposed action will have no effect on the population status of the Mexican vine snake.

Osprey (*Pandion haliaetus*)

This raptor is dark brown on its back and white on the underparts with a prominent dark eye stripe. The total range for the osprey is from Alaska to Newfoundland, along the Atlantic and Pacific coastlines, and in the Rocky Mountains south through central and South America. Within Arizona, the osprey occurs primarily in the White Mountains, along the Mogollon Rim, and along the Salt and Verde rivers. In southeastern Arizona, this raptor is an uncommon spring and fall transient, usually seen at ponds and reservoirs. Nesting habitat of the osprey consists of coniferous trees along rivers and lakes at elevations ranging from 1,829 to 2,377 m (6,000 – 7,800 ft) (AGFD 1997d).

Osprey population trends in Arizona are not well known. Only about 20 nest sites are known in the southwest, all within Arizona. This raptor is threatened by loss of nesting habitat and foraging perch sites. It is also threatened by recreational use of nesting habitat, shooting, and pesticide poisoning on wintering grounds (AGFD 1997d).

Because no construction will occur in perennial aquatic habitats, the proposed action will have no effect on the population status of the osprey.

Rose-throated becard (*Pachyramphus aglaiae*)

The rose-throated becard is a big-headed, thick billed bird that breeds in southeast Arizona, southern Texas (rare visitor along the Rio Grande), south through Mexico to Costa Rica. This species winters from northern Mexico south through to its breeding range. Within Arizona, rose-throated becards have been found breeding along Sonoita and Arivaca creeks, Sycamore Canyon (Atascosa Mountains), and Patagonia. Historically, this species nested in Guadalupe Canyon (east of Douglas) and near Tucson. Rose-throated becards typically inhabit marshes of Sonoran desertscrub communities of open to dense vegetation of shrubs, low trees, and succulents dominated by paloverde, prickly pear, and saguaro. This species also is found in the desert riparian deciduous woodland communities of marsh-woodlands, especially of cottonwoods, that occur where desert streams provide sufficient moisture for a narrow band of deciduous trees and shrubs along the margins. In Arizona, the rose-throated becard is found at elevations ranging from 1,082 to 1,228 m (3,550 – 4,030 ft) (AGFD 2001g).

Population trends for the rose-throated becard are currently unknown. Potential threats to this species include disturbance from bird watchers and degradation and loss of native riparian habitat through overgrazing, urban development, and groundwater depletion (AGFD 2001g). Because known occurrences of this species are outside the project area, the proposed action will have no effect on the population status of the rose-throated becard.

Thick-billed kingbird (*Tyrannus crassirostris*)

The thick-billed kingbird is a relatively stocky flycatcher with a large head and heavy bill. This kingbird occurs from southeastern Arizona and southwestern New Mexico south through western Mexico to western Guatemala. In Arizona, thick-billed kingbirds are most often seen around Sonoita and Arivaca creeks and in Madera and Guadalupe canyons. This species may occur in mountains of Pima, Santa Cruz and Cochise counties

where there are drainages with well-developed riparian areas. Habitat for the thick-billed kingbird consists of broad-leaved, riparian forests usually with well-developed large sycamores and cottonwoods at elevations ranging from 914 to 1,981 m (3,000 – 6,500 ft) (Tibbitts 1991).

Present distribution of the thick-billed kingbirds in Arizona is very limited. Potential threats include human recreational activities, encroachment of human development into breeding habitat, woodcutting, grazing, and groundwater depletion (Tibbitts 1991). Because no potential habitat occurs within the project area, the proposed action will have no effect on the population status of the thick-billed kingbird.

Tropical Kingbird (*Tyrannus melancholicus*)

The tropical kingbird is a large tyrant-flycatcher with a large bill and long, slightly notched tail. The tropical kingbird ranges from southeastern Arizona through western and central Mexico to central Argentina. Breeding birds have been found in Tucson, along the Santa Cruz Valley from Green Valley south, east of Phoenix in the Salt River Valley, to the San Pedro Valley. This species also has been reported from Sopori Wash. The Tropical Kingbird inhabits open and semi-open areas with scattered trees and shrubs. Also found in urban areas and roadsides with tall human-made fixtures (Stouffer and Chessser 1998).

Tropical kingbirds seem to persist or even thrive in developed areas. No negative effects of human activities have been reported (Stouffer and Chessser 1998). The proposed transmission line may cross potential habitat for this species; however, construction within riparian habitats will be minimized to the greatest extent possible. Placement of the transmission line may impact individual tropical kingbirds, however because of the linear nature of the project, only a small percentage of the population within the project area may be impacted. Furthermore, populations of this species occur outside of the project area. Therefore, impacts to tropical kingbirds are not likely to result in a trend toward federal listing or loss of viability.

6.0 LITERATURE CITED

- Abbate, D. J., A. Ditty, W. S. Richardson, and R. Olding. 1996. Cactus ferruginous pygmy-owl surveys and nest monitoring in the Tucson Basin area, Arizona. Final Report Internal Enhancement #U95503. Arizona Game and Fish Department, Phoenix, Arizona.
- _____, W. S. Richardson, R. L. Wilcox, M. J. Terrio, and S. M. Belhumeur. 1999. Cactus ferruginous pygmy-owl investigations in Pima and Pinal Counties, Arizona: 1997-1998. Region V Wildlife Program. Arizona Game and Fish Department, Tucson, AZ. 83 pp.
- _____, _____, _____, and S. Lantz. 2000. Cactus ferruginous pygmy-owl investigations in Pima and Pinal Counties, Arizona: 1999. Region V Wildlife Program. Arizona Game and Fish Department, Tucson, AZ. 37 pp.
- Aiken, C. E. H. 1937. (E.R. Warren, Ed.). Birds of the Southwest. Colorado College Publication, General Series No. 212:1-73.
- Alcorn, S. M., S. E. McGregor and G. Olin. 1962. Pollination requirements of the organpipe cactus. Cactus and Succulent Journal 34(5): 135-138.
- Allison, L. J., J. Rourke, and T. McCarthy. 2000. Characterizing vegetation in nesting and non-nesting plots for southwestern willow flycatchers within the same patch of continuous riparian vegetation. Page 15 in Abstracts of presented papers and posters, The Ecology and Conservation of the Willow Flycatcher, 24-26 October 2000, Arizona State University, Tempe, AZ.
- Arizona Game and Fish Department (AGFD). 1988. Threatened native wildlife in Arizona. Arizona Game and Fish Department, Phoenix, AZ. 32 pp.
- _____. 1991a. Chiltepine (*Capsicum annuum* var *glabriusculum*). Unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. 7 pp.
- _____. 1991b. Mexican vine snake (*Oxibelis aeneus*). Draft unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. 2 pp.
- _____. 1992. Sweet acacia (*Acacia smallii*). Draft unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. 4 pp.
- _____. 1993. Arizona wildlife views – bats of Arizona. Volume 36, Number 8. 36 pp.

- _____. 1994. Mexican long-tongued bat (*Choeronycteris mexicana*). Draft unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. 3 pp.
- _____. 1995a. Western barking frog (*Eleutherodactylus augusti cactorum*). Unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. 3 pp.
- _____. 1995b. Great Plains narrow-mouthed toad (*Gastrophryne olivacea*). Draft unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. 3 pp.
- _____. 1997a. Bartram's stonecrop (*Graptopetalum beartramii*). Unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. 8 pp.
- _____. 1997b. Pima Indian mallow (*Abutilon parishii*). Unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. 7 pp.
- _____. 1997c. Lowland leopard frog (*Rana yavapaiensis*). Unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. 3 pp.
- _____. 1997d. Cave myotis (*Myotis velifer*). Unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. 4 pp.
- _____. 1997e. Osprey (*Pandion haliaetus*). Draft unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. 4 pp.
- _____. 1998a. Beardless chinch weed (*Pectis imberbis*). Unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. 7 pp.
- _____. 1998b. Gentry indigo bush (*Dalea tentaculoides*). Unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. 8 pp.
- _____. 1998c. Large-flowered blue star (*Amsonia grandiflora*). Unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. 9 pp.

- _____. 1998d. Santa Cruz beehive cactus (*Coryphantha recurvata*). Draft unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. 5 pp.
- _____. 1998e. Santa Cruz striped agave (*Agave parviflora* ssp. *parviflora*). Draft unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. 4 pp.
- _____. 1998f. American peregrine falcon (*Falco peregrinus anatum*). Unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. 4 pp.
- _____. 1998g. Western yellow-billed cuckoo (*Coccyzus americanus occidentalis*). Unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. 3 pp.
- _____. 1998h. Southern pocket gopher (*Thomomys umbrinus intermedius*). Draft unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. 4 pp.
- _____. 1999a. Alamos deer vetch (*Lotus alamosanus*). Draft unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. 6 pp.
- _____. 1999b. Catalina beardtongue (*Penstemon discolor*). Draft unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. 8 pp.
- _____. 1999c. Chihuahuan sedge (*Carex chihuahuensis*). Draft unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. 6 pp.
- _____. 1999d. Chiricahua Mountain brookweed (*Samolus vagans*). Unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. 6 pp.
- _____. 1999e. Nodding blue-eyed grass (*Sisyrinchium cernuum*). Draft unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. 5 pp.
- _____. 1999f. Santa Cruz star leaf (*Choisya mollis*). Draft unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. 8 pp.

- _____. 1999g. Supine bean (*Macroptilium supinum*). Draft unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. 8 pp.
- _____. 1999h. Thurber hoary pea (*Tephrosia thurberi*). Draft unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. 6 pp.
- _____. 1999i. Virlet paspalum (*Paspalum virletii*). Draft unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. 3 pp.
- _____. 1999j. Weeping muhly (*Muhlenbergia xerophila*). Draft unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. 6 pp.
- _____. 1999k. Woolly fleabane (*Laennecia eriophylla*). Unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. 6 pp.
- _____. 2000a. Arid throne fleabane (*Erigeron arisolis*). Unpublished draft abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. 6 pp.
- _____. 2000b. Arizona giant sedge (*Carex ultra*). Unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. 8 pp.
- _____. 2000c. Broadleaf ground cherry (*Physalis latiphysa*). Draft Unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. 7 pp.
- _____. 2000d. Foetid passionflower (*Passiflora foetida*). Draft Unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. 7 pp.
- _____. 2000e. Lumholtz nightshade (*Solanum lumholtzianum*). Draft unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. 6 pp.
- _____. 2000f. Mock-pennyroyal (*Hedeoma dentatum*). Unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. 4 pp.

- _____. 2000g. Seeman groundsel (*Senecio carlomasonii*). Draft unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. 3 pp.
- _____. 2000h. Sonoran noseburn (*Tragia laciniata*). Draft unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. 7 pp.
- _____. 2000i. Superb beardtongue (*Penstemon superbus*). Draft unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. 3 pp.
- _____. 2000j. Thurber's morning glory (*Ipomoea thurberi*). Draft unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. 6 pp.
- _____. 2000k. Wiggins milkweed vine (*Metastelma mexicanum*). Unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. 6 pp.
- _____. 2000l. Northern gray hawk (*Asturina nitida maxima*). Draft unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. 5 pp.
- _____. 2001a. Arizona metalmark (*Calephelis rawsoni arizonensis*). Unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. 3 pp.
- _____. 2001b. Giant spotted whiptail (*Cnemidophorus burti strictogrammus*). Unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. 4 pp.
- _____. 2001c. Mexican garter snake (*Thamnophis eques megalops*). Unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. 4 pp.
- _____. 2001d. California leaf-nosed bat (*Macrotus californicus*). Unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. 5 pp.
- _____. 2001e. Sonoran Desert tortoise (*Gopherus agassizii*). Unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. 8 pp.

- _____. 2001f. Elegant trogon (*Trogon elegans*). Draft unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. 7 pp.
- _____. 2001h. Rose-throated becard (*Pachyramphus aglaiae*). Draft unpublished abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. 7 pp.
- _____. 2002. Loggerhead shrike (*Lanius ludovicianus*). Draft unpublished animal abstract compiled and edited by the Heritage Data Management System, Arizona Game and Fish Department, Phoenix, AZ. 4 pp.
- Arizona Interagency Desert Tortoise Team. 1996. Management plan for the Sonoran Desert population of the desert tortoise in Arizona.
- Avian Power Line Interaction Committee (APLIC). 1996. Suggested practices for raptor protection on power lines: the state of the art in 1996. Edison Electric Institute and Raptor Research Foundation. Washington, D.C.
- Bagley, B. E., D. A. Hendrickson, F. J. Abarca, and S. Hart. 1991. Status of the Sonoran topminnow (*Poeciliopsis occidentalis*) and desert pupfish (*Cyprinodon macularius*) in Arizona. Special Report on Project E5-2, Job 9, Title VI of the Endangered Species Act. Arizona Game and Fish Department, Phoenix, AZ.
- Bahre, C. J. 1991. A legacy of change. Historic human impact on vegetation in the Arizona borderlands. University of Arizona Press, Tucson, AZ.
- Baird, S. F. and C. Girard. 1853. Description of new species of fishes collected by Mr. John H. Clark, on the U.S.-Mexican boundary survey, under Lt. Col. Jas. D. Graham. Proceedings of the Academy of Natural Sciences of Philadelphia 6: 387-390.
- Bartlett, P. and R. Bartlett. 1999. A field guide to Texas reptiles and amphibians. Gulf Publishing Company, Houston, TX.
- Belnap, J. 1992. Potential role of cryptobiotic soil crusts in semiarid rangelands. Paper presented at the Symposium on Ecology, Management, and Restoration of Intermountain Annual Rangelands, Boise, ID, May 18-22, 1992.
- Belsky, A. J., and D. M. Blumenthal. 1997. Effects of livestock grazing on stand dynamics and soils in upland forests of the interior west. Conservation Biology 11(2): 315-327.

- Bendire, C. E. 1892. *Glaucidium phalaenoides* (Daudain), Ferruginous pygmy owl. Pages 409-411 *In* Life Histories of North American Birds with special reference to their habits and eggs. Smithsonian Institute, U.S. National Museum Special Bulletin No. 1.
- Bevanger, K. 1994. Bird interactions with utility structures: collision and electrocution, causes and mitigating measures. *Ibis* 54: 419-431.
- Blackburn, W. H. 1984. Impacts of grazing intensity and specialized grazing systems on watershed characteristics and responses. Pages 927-983 *In* Developing strategies for rangeland management. National Research Council/National Academy of Sciences. Westview Press, Boulder, CO.
- Bockstanz, L. 1998. Herps of Texas – lizards: Texas Horned Lizard (*Phrynosoma cornutum*).
<http://www.zo.utexas.edu/research/txherps/lizards/phrynosoma.cornutum.html>.
22 September 1999.
- Brooks, J. E. 1985. Factors affecting the success of Gila topminnow introductions on four Arizona National Forests. Report to U.S. Fish and Wildlife Service, Office of Endangered Species, Albuquerque, New Mexico. Arizona Game and Fish Department, Phoenix, AZ.
- _____. 1986. Status of natural and introduced Sonoran topminnow (*Poeciliopsis o. occidentalis*) populations in Arizona through 1985. Report to U.S. Fish and Wildlife Service, Office of Endangered Species, Albuquerque, New Mexico. Arizona Game and Fish Department, Phoenix, AZ.
- Brown, D. E., C. H. Lowe, and J. F. Hausler. 1977. Southwestern riparian communities: their biotic importance and management in Arizona *In* R. R. Johnson and D. A. Jones [Eds.], Importance, preservation, and management of riparian habitats: a symposium. General Technical Report RM-43. USDA Forest Service, Denver, CO.
- _____. 1983a. On the status of the jaguar in the Southwest. *Southwestern Naturalist* 28: 459-460.
- _____. 1983b. The wolf in the southwest. University of Arizona Press, Tucson, AZ. 195 pp.
- _____. 1985. Black-bellied whistling duck. Pages 110 – 112 *In* Arizona Wetlands and Waterfowl. The University Press, Tucson, AZ.
- _____. 1991. Revival for El Tigre? *Defenders* 66(1): 27-35.

- _____. (Ed.). 1994. Biotic communities: southwestern United States and northwestern Mexico. University of Utah Press, Salt Lake City, UT. 342 pp.
- Brown, M. and F. J. Abarca. 1992. An update status report of the Sonoran topminnow, *Poeciliopsis occidentalis*, and desert pupfish, *Cyprinodon macularius*, in Arizona. Arizona Game and Fish Department, Phoenix.
- Browning, M. R. 1993. Comments on the taxonomy of *Empidonax traillii* (willow flycatcher). Western Birds 24: 241-257.
- Busby, F. E. and G. F. Gifford. 1981. Effects of livestock grazing on infiltration and erosion rates measured on chained and unchained pinyon-juniper sites in southeastern Utah. Journal of Range Management 34: 400-405.
- Cartron, J. E., W. S. Richardson, and G. A. Proudfoot. 2000a. The cactus ferruginous pygmy-owl: taxonomy, distribution, and natural history. Pages 5-15 In J. E. Cartron and D. M. Finch [Eds.], Ecology and conservation of the cactus ferruginous pygmy-owl in Arizona. Gen. Tech. Rpt. RMRS-GTR-43. USDA Forest Service, Rocky Mountain Research Station, Ogden, UT. 68 pp.
- Cartron, J. E., S. H. Soleson, S. Russell, G. A. Proudfoot, and W. S. Richardson. 2000b. The ferruginous pygmy-owl in the tropics and at the northern end of its range: habitat relationships and requirements. Pages 47-53 In J. E. Cartron and D. M. Finch [Eds.], Ecology and conservation of the cactus ferruginous pygmy-owl in Arizona. Gen. Tech. Rpt. RMRS-GTR-43. USDA Forest Service, Rocky Mountain Research Station, Ogden, UT.
- Cockrum, E.L. 1991. Seasonal distribution of northwestern populations of the long-nosed bat, *Leptonycteris sanborni*, Family Phyllostomidae. Anales Inst. Biol. Univ. Nac. Auton. Mexico, Ser. Zool. 62(2): 181-202.
- Constantine, D.G. 1966. New bat locality records from Oaxaca, Arizona and Colorado. Journal of Mammalogy 47(1): 125-126.
- _____. 1998. Range extensions of ten species of bats in California. Bulletin of Southern California Academy of Science 97(2): 49-75.
- Constantz, G. D. 1974. Reproductive effort in *Poeciliopsis occidentalis* (Poeciliidae). Southwestern Naturalist 19: 47-52.
- _____. 1975. Behavioral ecology of mating in the male Gila topminnow, *Poeciliopsis occidentalis* (Cyprinodontiformes: Poeciliidae). Ecology 56: 966-973.
- _____. 1976. Life history strategy of the Gila topminnow, *Poeciliopsis occidentalis*: a field evaluation of theory on the evolution of life histories. Ph.D. Dissertation, Arizona State University, Tempe.

- _____. 1979. Life history patterns of a livebearing fish in contrasting environments. *Oecologia* 40: 189-201.
- _____. 1989. Reproductive biology of poeciliid fishes. Pages 33-50 *In* G. K. Meffe and F. F. Snelson Jr. [Eds.], *Ecology and Evolution of Livebearing Fishes (Poeciliidae)*. Prentice Hall, Englewood Cliffs.
- Corman, T. E. 1994. Black-bellied whistling duck. Page 18 *In* Nongame Field Notes. AGFD Wildlife News, Arizona.
- Coronado National Forest (CNF). 2000. FY 1999 Annual monitoring and evaluation report. Coronado National Forest report, 27 pp.
- Coues, E. 1872. A new bird to the United States. *American Naturalist* 6: 370.
- Dalton, V. M. and D. C. Dalton. 1993. Assessment of the impacts of low-level military aircraft on *Leptonycteris curasoae*, an endangered bat, at Organpipe Cactus National Monument, Arizona. Contract Nos. USAF/Luke AFB F0260492MS359 and NPS/ORPI O.M. 1443 PX866092081. Report to Organpipe Cactus National Monument and 58th Civil Eng. Sq. Luke AFB. 54 pp.
- Dalton, V. M., D. C. Dalton and S. L. Schmidt. 1994. Roosting and foraging use of a proposed military training site by the long-nosed bat, *Leptonycteris curasoae*. Contract Nos. DACA65-94-M-0831 and DACA65-94-M-0753. Report to Luke Air Force Natural Resources Program. 34 pp.
- DeBano, L. F. and L. J. Schmidt. 1989. Interrelationships between watershed condition and health of riparian areas of southwestern United States. Pages 45-52 *In* M. T. Billings, R. E. Gresswell, B. A. Barton, and J. L. Kershner [Eds.], *Practical approaches to riparian resource management. An educational workshop*. May 8-11, 1989. U.S. Bureau of Land Management, Billings, MT.
- Desert Tortoise Council. 1994 (Revised 1999). Guidelines for handling desert tortoises during construction projects. Edward L LaRue, Jr. [Ed.]. Wrightwood, CA. 19 pp.
- Drost, C. A. 1998. Survey of amphibians and amphibian habitat on the upper Santa Cruz River, Arizona. USGS, Biological Resources Division/Colorado Plateau Research Station, Flagstaff, AZ.
- _____, M. K. Sogge, and E. Paxton. 1998. Preliminary diet study of the endangered southwestern willow flycatcher. Report to U. S. Bureau of Reclamation. USGS, Biological Resources Division/Colorado Plateau Research Station, Flagstaff, AZ. 26 pp.

- Fleming, T. H. 1991. Nectar corridors and the diet of migratory nectarivorous bats. *Bat Research News* 33: 71.
- _____. 1994. Lesser long-nosed bat recovery plan. U. S. Fish and Wildlife Service, Albuquerque, NM. 49 pp.
- _____. 1995. Lesser long-nosed bat recovery plan. U.S. Fish and Wildlife Service. Albuquerque, NM. Page 29.
- Forrest, R. E. 1992. Habitat use and preference of Gila topminnow. M.S. Thesis, University of Arizona, Tucson, AZ.
- Gentry, H. S. 1982. Agaves of Continental North America. University of Arizona Press, Tucson, AZ. Pages 443-447 and 538-545.
- Gerking, S. D. and D. V. J. Plantz. 1980. Size-biased predation by the Gila topminnow, *Poeciliopsis occidentalis* (Baird and Girard). *Hydrobiologia* 72: 179-191.
- Gifford, G. F. and R.H. Hawkins. 1978. Hydrologic impact of grazing on infiltration: a critical review. *Water Resour. Res.* 14: 305-313.
- Glenn, W. 1996. Eyes of fire: encounter with a borderlands jaguar. Printing Corner Press, El Paso, TX.
- Glinski, R. L. 1998. The raptors of Arizona. Arizona Game and Fish Department and University of Arizona Press, Tucson, AZ. 220 pp.
- Goldman, E. A. 1932. The jaguars of North America. *Proc. Biol. Soc. Washington* 45: 143-146.
- Green, L. R. 1977. Fuelbreaks and other fuel modifications for wildland fire control. *Agric. Handb.* 499. Washington, DC: U.S. Department of Agriculture, Forest Service. 79 pp.
- Greenberg, C. H.; Crownover, S. H.; Gordon, D. R. 1997. Roadside soils: a corridor for invasion of xeric scrub by nonindigenous plants. *Natural Areas Journal*. 17(2): 99-109.
- Gucinski, H.; M. J. Furniss, R. R. Ziemer, and M. H. Brookes. 2001. Forest roads: a synthesis of scientific information. Gen. Tech. Rep. PNWGTR-509. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 103 pp.
- Hall, E. R. 1981. The mammals of North America. 2nd Edition. John Wiley and Sons, New York, 2: 601-1181 + 90.

- Harness, R. E. and K. R. Wilson. 2000. Raptor electrocutions and outages: a review of rural utility records spanning 1986-1996. Pages 765-771 *In* R. D. Chancellor and B. U. Meyburg [Eds.], *Raptors at risk*. Hancock House Publishers, Blaine, WA.
- Harris, L. K. and R. B. Duncan. 1998. Cactus ferruginous pygmy-owl 1998 survey of southern Arizona, final report. United States Fish and Wildlife Service grant agreement # 1448-20181-98-G915. 39 pp.
- _____. and _____. 1999. Pima County cactus ferruginous pygmy-owl survey, final report. Pima County Government Contract #07-26-H-125828-0499. 78 pp.
- Harvey, M. J., J. S. Altenbach, and T. L. Best. 1999. Bats of the United States. Arkansas Game and Fish Commission and U.S. Fish and Wildlife Service – Asheville Field Office. 64 pp.
- Hatten, J., C. Paradzick, and M. Sumner. 2000. Constructing a nest-site suitability model for the southwestern willow flycatcher in Arizona. Page 20 *In* Abstracts of presented papers and posters, *The Ecology and Conservation of the Willow Flycatcher*, 24-26 October 2000, Arizona State University, Tempe.
- Hayward, B. and E. L. Cockrum. 1971. The natural history of the western long-nosed bat, *Leptonycteris sanborni*. Western New Mexico University, Research Science 1(2): 74-123.
- Heath, W. G. 1962. Maximum temperature tolerance as a function of constant temperature acclimation in the Gila topminnow, *Poeciliopsis occidentalis*. Ph.D. Dissertation, University of Arizona, Tucson.
- Hoffmeister, D. F. 1986. *Mammals of Arizona*. University of Arizona Press, Tucson, AZ. 602 pp.
- Howell, D. J. and B. S. Roth. 1981. Sexual reproduction in agaves: the benefits of bats; cost of semelparous advertising. *Ecology* 62: 3-7.
- Howell, S. N. G. and S. Webb. 1995. *A guide to the birds of Mexico and northern Central America*. Oxford University Press, New York, NY. 851 pp.
- Hubbs, C. L. and R. R. Miller. 1941. Studies of the fishes of the order Cyprinodontes. XVII – genera and species of the Colorado River system. *Occasional Papers of the Museum of Zoology, University of Michigan* 433: 1-9.
- Hubbard, J. P. 1987. The status of the willow flycatcher in New Mexico. *Endangered Species Program*, New Mexico Department of Game and Fish, Santa Fe, NM. 29 pp.

- Hughes, J. M. 1999. Yellow-billed cuckoo (*Coccyzus americanus*). In *The Birds of North America*, No. 418 (A. Poole and F Gill, eds.). The Birds of North America, Inc., Philadelphia, PA. 28 pp.
- Jennings, M. R. 1987. Status of the western painted turtle (*Chrysemys picta bellii*) in Arizona. *Journal of the Arizona-Nevada Academy of Sciences* 22: 129-133.
- Johnson, R. R. and L.T. Haight. 1985. Avian use of xeroriparian ecosystems in the North American warm deserts. Pages 156-160 In R. R. Johnson et al., technical coordinator. *Riparian ecosystems and their management: reconciling conflicting uses*.
- Kearney, T. H. and R. H. Peebles. 1960. Arizona flora – 2nd edition. University of California Press, Berkeley and Los Angeles, CA. 1,085 pp.
- Lange, K. K. 1960. The jaguar in Arizona. *Transactions of the Kansas Academy of Science* 63: 96-101.
- Lesh, T. D. and T. E. Corman. 1995. Cactus ferruginous pygmy-owl surveys in Arizona: 1993-1995. Nongame and Endangered Wildlife Program Technical Report 76. Arizona Game and Fish Department, Phoenix, AZ. 23 pp.
- Lonsdale, W. M and A. M. Lane. 1994. Tourist vehicles as vectors of weed seeds in Kadudu National Park, northern Australia. *Biological Conservation* 69(3): 277-283.
- Lyon, L. J. 1983. Road density models describing habitat effectiveness for elk. *Journal of Forestry*. 81: 592-595.
- Marsh, P. C. and W. L. Minckley. 1990. Management of endangered Sonoran topminnow at Bylas Springs, Arizona. Description, critique and recommendations. *Great Basin Naturalist* 50(3): 265-272.
- Marti, C. D. 2002. Enhancing raptor populations, a technique manual. The Peregrine Fund. Boise, ID. 77 pp.
- Maynard, W. R. 1995. Summary of 1994 survey effort in New Mexico for southwestern willow flycatcher (*Empidonax traillii extimus*). Contract #94-516-69. New Mexico Department of Game and Fish, Santa Fe, New Mexico. 48 pp.
- McCarthy, T. D., C. E. Paradzick, J. W. Rourke, M. W. Sumner, and R. F. Davidson. 1998. Arizona partners in flight southwestern willow flycatcher 1997 survey and nest monitoring report. Report to the Bureau of Reclamation, Phoenix, AZ.
- McGregor, S. E., S. M. Alcorn, and G. Olin. 1962. Pollination and pollinating agents of the saguaro. *Ecology* 43: 259-267.

- McNeil, R., J. R. Rodriguez Sr., and H. Ouellet. 1985. Bird mortality at a power transmission line in northeastern Venezuela. *Biological Conservation* 31: 151-165.
- McPherson, G. R. 1995. The role of fire in desert grasslands. Pages 130-151 *In* M. P. McClaran and T. R. Van Devender [Eds.], *The Desert Grassland*. University of Arizona Press, Tucson, AZ.
- Mech, L. D. 1973. Wolf numbers in the Superior National Forest of Minnesota. Res. Pap. NC-97. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Research Experiment Station. 10 pp.
- Meffe, G. K., D. A. Hendrickson, and W. L. Minckley. 1983. Factors resulting in decline of the endangered Sonoran topminnow, *Poeciliopsis occidentalis* (Antheriniformes: Poeciliidae) in the United States. *Biological Conservation* 25:135-159.
- _____. 1984. Effects of abiotic disturbance on coexistence of predator-prey fish species. *Ecology* 65(5): 1525-1534.
- Minckley, W. L. 1973. *Fishes of Arizona*. Arizona Game and Fish Department, Sims Printing Company, Inc., Phoenix, AZ. 293 pp.
- _____. and J. E. Brooks. 1985. Transplantations of native Arizona fishes: records through 1980. *Journal of the Arizona-Nevada Academy of Sciences* 20(2): 73-90.
- Mladenoff, D. J., T. A. Sickley, R. G. Haight, and A. P. Wydevens. 1995. A regional landscape analysis and prediction of favorable gray wolf habitat in the northern Great Lakes region. *Conservation Biology* 9(2): 279-294.
- Monson, G. and A. R. Philips. 1981. Annotated checklist of the birds of Arizona, 2nd edition, University of Arizona Press, Tucson, AZ. 292 pp.
- _____. 1987. An "almost certain" ferruginous pygmy-owl was heard along lower Aravaipa Canyon (Woods Ranch now owned by TNC) on 28 July 1987. *American Birds* 41:1471.
- _____. 1998. Ferruginous pygmy-owl. Pages 159-161 *In* R. Glinski [Ed.]. *The Raptors of Arizona*. University of Arizona Press, Tucson, AZ. 220 pp.
- Morrison, J. L. 1996. Crested Caracara (*Caracara plancus*) *In* *The Birds of North America*, No. 249 (A. Poole and F. Gill, eds.). The Academy of Natural Sciences, Philadelphia, PA, and The American Ornithologists' Union, Washington, DC.

- Muiznieks, B. D., S. J. Sferra, T. E. Corman, M. K. Sogge, and T. J. Tibbitts. 1994. Arizona Partners in Flight southwestern willow flycatcher survey 1993. Draft report: Nongame and endangered wildlife program, Arizona Game and Fish Department, Phoenix, AZ. Draft of April 1994. 28 pp.
- Nelson, E.W. and E.A. Goldman. 1933. Revision of the jaguars. *Journal of Mammalogy* 14:221-240.
- New Mexico Game and Fish Department and The Fish and Wildlife Information Exchange. 2000. Five-Stripped Sparrow (*Aimophila quinquestriata*). [Web page] No archive known [Accessed 7 June 2002]. BISON. http://www.fw.vt.edu/fishex/nmex_main/species/041830.htm
- Nobel, T. A. 1995. Birds and power lines: selected interaction and management issues in the electric utility industry. M. S. Thesis, Prescott College, Prescott, AZ.
- Nowak, R. M. 1994. Walker's bats of the world. Johns Hopkins University Press, Baltimore, MD. 287 pp.
- Paradzick, C. E., R. F. Davidson, J. W. Rourke, M. W. Sumner, and T. D. McCarthey. 1999. Southwestern willow flycatcher 1998 survey and nest monitoring report. Nongame and Endangered Wildlife Program, Arizona Game and Fish Department, Phoenix, AZ, Technical Report 141.
- _____, _____, _____, _____, A. M. Wartell, and T. D. McCarthey. 2000. Southwestern willow flycatcher 1999 survey and nest monitoring report. Nongame and Endangered Wildlife Program Technical Report 151. Arizona Game and Fish Department, Phoenix, AZ. 93 pp.
- Parsons, D. R. 1996. Case study: the Mexican wolf. Pages 101-123 *In* E. A. Herrera and L. F. Huenneke [Eds.] *New Mexico's Natural Heritage: Biological Diversity in the Land of Enchantment*. New Mexico Journal of Science, Volume 36.
- Peterson, R. T. 1990. A field guide to western birds. Third Edition. Houghton Mifflin Company, Boston, MA. 432 pp.
- Phillips, A., J. Marshall, and G. Monson. 1983. The birds of Arizona. University of Arizona Press, Tucson, AZ. 212 pp.
- Phillips, A. R. 1948. Geographic variation in *Epidonax traillii*. *The Auk* 65: 507-514.
- _____, J.T. Marshall Jr., and G. Monson. 1964. The Birds of Arizona. University of Arizona Press, Tucson, AZ. 220 pp.

- Pima County. 2001. Priority vulnerable species: analysis and review of species proposed for coverage by the multiple-species conservation plan DRAFT. Pima County, AZ. 385 pp.
- Proudfoot, G. and S. M. Beasom. 1997. Food habits of nesting ferruginous pygmy-owls in southern Texas. *Wilson Bulletin* 109:741-748.
- _____. and R. D. Slack. 2001. Comparisons of ferruginous pygmy-owl mtDNA at local and international scales. Department of Wildlife and Fisheries Sciences, Texas A & M University. A Report for Pima County. 11 pp.
- Pyle, P. 1997. Identification guide to North American birds. Part I Columbidae to Ploceidae. Slate Creek Press, Bolinas, CA.
- Richardson, W. S., J. E. Cartron, D. J. Krueper, L. Turner, and T. H. Skinner. 2000. The status of the cactus ferruginous pygmy-owl in Arizona: Population surveys and habitat assessment. Pages 27-39 *In*: J. E. Cartron and D. M. Finch [Eds.], Ecology and conservation of the cactus ferruginous pygmy-owl in Arizona. General Technical Report, RMRS-GTR-43, Ogden, Utah. U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. 68 pp.
- Ridgely, R. S. and G. Tudor. 1994. The birds of South America: suboscine passerines. University of Texas Press, Austin, TX.
- Rogers, G. F. and J. Steele. 1980. Sonoran desert fire ecology. Pages 15 -16 *In* M. A. Stokes and J. H. Dieterich [Tech. Coord.]. Proceedings of the fire history workshop. United States Forest Service General Technical Report RM-81.
- Roller, P. S. 1996. Distribution, growth and reproduction of Pima pineapple cactus (*Coryphantha scheeri* Kuntz var. *robustispina* Schott). M. S. Thesis, University of Arizona.
- Rudis, V. A. 1995. Regional forest fragmentation effects on bottomland hardwood community types and resource values. *Landscape Ecology* 10(5): 291-307.
- San Diego Natural History Museum. 1995. *Empononax traillii extimus* in California. The willow flycatcher workshop. 17 November 1995. 66 pp.
- Salazar, L. A. and A. Gonzalez-Caban. 1987. Spatial relationship of a wildfire, fuelbreaks, and recently burned areas. *Western Journal of Applied Forestry* 2(2): 55-58.
- Schoenherr, A. A. 1974. Life history of the topminnow, *Poeciliopsis occidentalis* (Baird and Girard) in Arizona, and an analysis of its interaction with the mosquitofish *Gambusia affinis* (Baird and Girard). Ph.D. Dissertation, Arizona State University, Tempe.

- _____. 1977. Density dependent and density independent regulation of reproduction in the Gila topminnow, *Poeciliopsis occidentalis* (Baird and Girard). *Ecology* 58:438-444.
- Schultz, R. J. 1961. Reproductive mechanism of unisexual and bisexual strains of the viviparous fish *Poeciliopsis*. *Evolution* 15:302-325.
- Sferra, S. J., R. A. Meyer, and T. E. Corman. 1995. Arizona Partners in Flight 1994 southwestern willow flycatcher survey. Final Technical Report 69. Arizona Game and Fish Department, Nongame and Endangered Wildlife Program, Phoenix, AZ. 46 pp.
- _____, T. E. Corman, C. E. Paradzick, J. W. Rourke, J. A. Spencer, and M. W. Sumner. 1997. Arizona Partners in Flight southwestern willow flycatcher survey 1993-1996 summary report. Technical Report. Arizona Game and Fish Department, Nongame and Endangered Wildlife Program, Phoenix, AZ. 97 pp.
- Simms, J. K. and K. M. Simms. 1992. What constitutes quality habitat for Gila topminnow (*Poeciliopsis occidentalis*)? An overview of habitat parameters supporting a robust population at Cienega Creek, Pima County, AZ. Abstract. *Proceedings of the Desert Fishes Council* 22-23:82.
- Simons, L. H. 1987. Status of the Gila topminnow (*Poeciliopsis o. occidentalis*) in the United States. Special Report on Project E-1, Title VI of the Endangered Species Act of 1973, Arizona Game and Fish Department, Phoenix.
- Smith, A. B., C. E. Paradzick, A. A. Woodward, P. E. T. Dockens, and T. D. McCarthy. 2002. Southwestern willow flycatcher 2001 survey and nest monitoring report. Technical Report 191. Arizona Game and Fish Department, Nongame and Endangered Wildlife Program, Phoenix, AZ. 63 pp.
- Sogge, M. K. and T. J. Tibbitts. 1992. Southwestern willow flycatcher (*Empidonax traillii extimus*) surveys along the Colorado River in Grand Canyon National Park and Glen Canyon National Recreation Area. NPS CPSU/Northern Arizona University, Flagstaff, AZ. 43 pp.
- _____, _____, and S. J. Sferra. 1993. Status of the southwestern willow flycatcher along the Colorado River between Glen Canyon Dam and Lake Mead – 1993. Summary Report. National Park Service Cooperative Park Studies Unit/Northern Arizona University, U. S. Fish and Wildlife Service, and Arizona Game and Fish Department, Flagstaff, AZ. 69 pp.
- _____, and _____. 1994. Distribution and status of the southwestern willow flycatcher along the Colorado River in the Grand Canyon – 1994. Summary Report. National Biological Service, Colorado Plateau Research Station, Northern Arizona University, Flagstaff, AZ. 37 pp.

- _____, R. M. Marshall, S. J. Sferra, and T. J. Tibbitts. 1997. A southwestern willow flycatcher survey protocol and breeding ecology summary. National Park Service/Colorado Plateau Research Station, Northern Arizona University, Technical Report NRTR-97/12. 37 pp.
- Soule, M. E. 1986. Conservation biology. The science of scarcity and diversity. Sinauer Association, Inc. Sunderland, MA. 584 pp.
- Spencer, J. A., S. J. Sferra, T. E. Corman, J. W. Rourke, and M. W. Sumner. 1996. Arizona Partners in flight 1995 southwestern willow flycatcher survey. Technical Report 97, March 1996. Arizona Game and Fish Department, Phoenix, AZ. 69 pp.
- Spencer, J. A. and D. P. Humphrey. 1999. Species of concern at the Arizona Army National Guard's Florence Military Reservation, Pinal County, AZ. Arizona Army National Guard. 32 pp.
- State of Arizona. 1990. Final report and recommendations of the Governor's riparian habitat task force. Executive Order 89-16. Streams and riparian resources. October 1990. Phoenix, AZ. 28 pp.
- Stiles, F. G. and A. F. Skutch. 1989. A guide to the birds of Costa Rica. Comstock, Ithaca, New York. 364 pp.
- Stouffer, P. C. and R. T. Chesser. 1998. Tropical Kingbird (*Tyrannus melancholicus*) In A. Poole and F. Gill, [Eds.], The Birds of North America, No. 358. The Academy of Natural Sciences, Philadelphia, PA, and The American Ornithologists' Union, Washington, DC.
- Stromberg, J. C., J. A. Tress, J. D. Wilkins, and S. D. Clark. 1992. Response of velvet mesquite to groundwater decline. Journal of Arid Environments 23: 45-58.
- _____. 1993a. Fremont cottonwood-Goodding willow riparian forests: a review of their ecology, threats, and recovery potential. Journal of the Arizona-Nevada Academy of Science 26(3): 97-110.
- _____. 1993b. Fremont cottonwood-Goodding willow riparian forests: a review of their ecology, threats, and recovery potential. Journal of the Arizona-Nevada Academy of Science 27(1): 111-124.
- Swank, W. G. and J. G. Teer. 1989. Status of the jaguar-1987. Oryx 23: 14-21.
- Tetra Tech, Inc. 2003. Draft TEP Sahuarita - Nogales transmission line Environmental Impact Statement.

- Thiel, R. P. 1985. The relationship between road densities and wolf habitat suitability in Wisconsin. *American Midland Naturalist* 113(2): 404-407.
- Thomas, P. A. 1991. Response of Succulents to Fire: A Review. *International Journal of Wildland Fire* 1(1): 11 – 22.
- Tibbitts, T. J. 1991. Thick-billed kingbird (*Tyrannus crassirostris*). *Wildlife Views*, Arizona Game and Fish Department. Page 17.
- Unitt, P. 1987. *Empidonax traillii extimus*: An endangered subspecies. *Western Birds* 18: 137-162.
- URS. 2003. Tucson Electric Power Sahuarita-Nogales transmission line project - roads analysis.
- U. S. Census Bureau. 2000. Arizona state and county facts, deprived from population estimates, 2000 census of population and housing, 1990 census of population and housing, small area income and poverty estimates, county business patterns, and 1997 census of governments.
<http://quickfacts.census.gov/qfd/states/04/04023.html>.
- U.S. Department of the Interior. 1967. Native fish and wildlife. Endangered Species. *Federal Register* 32(48): 4001.
- U.S. Fish and Wildlife Service (USFWS). 1982. Mexican Wolf Recovery Plan. USFWS, Albuquerque, NM. 103 pp.
- _____. 1984. Gila and Yaqui Topminnow Recovery Plan. USFWS, Albuquerque, NM.
- _____. 1988. Endangered and threatened wildlife and plants; determination of endangered status for two long-nosed bats. *Federal Register* 53(190): 38456-38460.
- _____. 1990. Listed cats of Texas and Arizona recovery plan (with emphasis on the ocelot). U.S. Fish and Wildlife Service, Albuquerque, New Mexico. 131 pp.
- _____. 1993. Determination of endangered status for the plant Pima pineapple cactus (*Coryphantha scheeri* var. *robustispina*). *Federal Register* 58 (158): 49875-49880.
- _____. 1997a. Endangered and threatened wildlife and plants: determination of endangered status for the cactus ferruginous pygmy-owl in Arizona. *Federal Register* 62: 10730-10747.

- _____. 1997b. Final determination of critical habitat for the southwestern willow flycatcher. Federal Register 62 (140): 39129-39146.
- _____. 1999. Endangered and threatened wildlife and plants; Final Rule: designation of critical habitat for the cactus ferruginous pygmy-owl. Federal Registrar 64: 37419-37440.
- _____. 2000. Recommended guidance for private landowners concerning the cactus ferruginous pygmy-owl: and the cactus ferruginous pygmy-owl survey protocol. Federal Register 65: 14999-15000.
- _____. 2001a. Biological Opinion on the effects of the scour protection project on I-19 and Frontage Road bridges over Peck Canyon near Santa Cruz River confluence in Santa Cruz County, Arizona. 38 pp.
- _____. 2001b. Biological Opinion on the effects of the proposed construction of Madera Highlands east of Green Valley in Pima County, Arizona. 18 pp.
- _____. 2001c. Biological Opinion on the effects of proposed livestock grazing and management on the Montana Allotment on the Nogales Ranger District, Santa Cruz County, Arizona. 51 pp.
- _____. 2001d. Mexican gray wolf (*Canis lupus baileyi*). Unpublished abstract, U. S. Fish and Wildlife Service, Arizona Ecological Services. 1 page.
- _____. 2002a. Biological Opinion on the effects of the Thornydale Road improvement project in Pima County, AZ. 66 pp.
- _____. 2002b. Biological Opinion on the effects of the proposed Pueblo Oasis Development in Pima County, AZ. 60 pp.
- _____. 2002c. Biological Opinion on the effects of the proposed issuance of a NPDES permit for ASARCO's Mission Mine Complex in Pima County, AZ. 15 pp.
- U.S. Forest Service (USFS). 1990. Soil and Water Conservation Practices Handbook, Region 3 FSH2509.22
- U.S. Geological Survey (USGS). 2003. Northern Prairie Wildlife Research Center. North Dakota's federally listed endangered, threatened, and candidate species – 1995: Western burrowing owl (*Athene cunicularia hypugea*). <http://www.npwrc.usgs.gov/resource/distr/others/nddanger/species/athecuni.htm>. 28 March 2003.
- Weedman, D. A. and K. L. Young. 1997. Status of the Gila topminnow and desert pupfish in Arizona. Arizona Game and Fish Department, Phoenix, AZ.

Wilcox, R., W. S. Richardson, and D. Abbate. 2000. Habitat selection by cactus ferruginous pygmy-owls in southern Arizona: Preliminary results. Arizona Game and Fish Department, Phoenix, AZ. 43 pp.

7.0 LIST OF ACRONYMS

ACC	Arizona Corporation Commission
ADEQ	Arizona Department of Environmental Quality
AGFD	Arizona Game and Fish Department
AOU	American Ornithologists' Union
ASLD	Arizona State Land Department
AUM	Animal Unit per Month
BA	Biological Assessment
BLM	Bureau of Land Management
BMP	Best Management Practices
BO	Biological Opinion
CFPO	Cactus Ferruginous Pygmy Owl
Citizens	Citizens Communications
CNF	Coronado National Forest
DBH	Diameter Breast Height
DOE	Department of Energy
EMA	Ecosystem Management Area
ESA	Endangered Species Act
GPS	Global Positioning System
HDMS	Heritage Data Management System
HEG	Harris Environmental Group, Inc.
I-19	Interstate 19
LLNB	Lesser Long-nosed Bat
MOA	Memorandum of Agreement
NPDES	National Pollutant Discharge Elimination System
OHV	Off-Highway Vehicle
PPC	Pima Pineapple Cactus
RA	Roads Analysis

RD	Ranger District
RNA	Research Natural Area
ROW	Right-of-way
RU	Recovery Units
SL	Standard Length
SWFL	Southwestern Willow Flycatcher
TEP	Tucson Electric Power
UDI	Undocumented Immigrants
USDOI	United States Department of Interior
USFWS	United States Fish and Wildlife Service
USFS	United States Department of Agriculture Forest Service
YOY	Young-of-the-year

APPENDIX A

Natural Resource Agencies Correspondence.

1. U. S. Fish and Wildlife Service, dated 14 May 2002.
2. Arizona Game and Fish Department, dated 25 April 2002.

APPENDIX B

**Plants documented along proposed ROW of the TEP Citizens Interconnect Project,
July to October 2002.**

	SCIENTIFIC NAME	COMMON NAME	FAMILY
CACTUS & SUCCULENTS			
	<i>Agave parryi</i>	century plant	Agavaceae
	<i>Agave schottii</i>	shindagger	Agavaceae
	<i>Coryphantha scheeri</i> var. <i>robustispina</i>	Pima pineapple cactus	Cactaceae
	<i>Dasyliion wheeleri</i>	sotol	Agavaceae
	<i>Echinocereus</i> spp.	hedgehog cactus	Cactaceae
	<i>Echinocereus pectinatus</i> var. <i>rigidissimus</i>	Arizona rainbow cactus	Cactaceae
	<i>Ferocactus wislizenii</i>	fishhook barrel cactus	Cactaceae
	<i>Fouquieria splendens</i>	ocotillo	Fouquieriaceae
	<i>Mammillaria</i> spp.	pincushion cactus	Cactaceae
	<i>Nolina microcarpa</i>	beargrass	Agavaceae
	<i>Opuntia</i> spp.	cholla	Cactaceae
	<i>Opuntia</i> spp.	prickly pear	Cactaceae
	<i>Opuntia spinosior</i>	walkingstick cactus	Cactaceae
	<i>Yucca elata</i>	soaptree yucca	Agavaceae
GRASSES			
	<i>Bouteloua barbata</i> or <i>B. rothrockii</i>	six-weeks or Rothrock grama	Poaceae
	<i>Bothriochloa barbinodis</i>	cane beard grass	Poaceae
	<i>Bouteloua curtipendula</i>	side oats grama	Poaceae
	<i>Bouteloua gracilis</i>	blue grama	Poaceae
	<i>Bouteloua hirsuta</i>	hairy grama	Poaceae
	<i>Bouteloua parryi</i>	Parry grama	Poaceae
	<i>Bouteloua repens</i>	slender grama	Poaceae
	<i>Digitaria californica</i>	Arizona cottontop	Poaceae
	<i>Erioneuron pulchellum</i>	fluffgrass	Poaceae
	<i>Hilaria belangeri</i>	curly mesquite	Poaceae
	<i>Leptochloa dubia</i>	green sprangletop	Poaceae
	<i>Muhlenbergia emersleyi</i>	bull grass	Poaceae
	<i>Muhlenbergia rigens</i>	deer grass	Poaceae
	<i>Piptochaetium fimbriatum</i>	pinyon rice grass	Poaceae
	<i>Sporobolus</i> spp.	dropseed	Poaceae
FORBS			
	<i>Abutilon incanum</i>	Indian mallow	Malvaceae
	<i>Allionia incarnata</i>	trailing windmills	Nyctaginaceae
	<i>Ambrosia confertiflora</i>	weakleaf burr ragweed	Asteraceae
	<i>Amoreuxia palmatiflida</i>	Arizona yellow show	Cochlospermaceae
	<i>Argemone</i> sp.	prickly poppy	Papaveraceae
	<i>Artemisia ludoviciana</i>		Asteraceae
	<i>Asclepias asperula</i>	antelope horns	Asclepiadaceae
	<i>Asclepias nummularia</i>	tufted milkweed	Asclepiadaceae
	<i>Asclepias tuberosa</i>	butterfly milkweed	Asclepiadaceae
	<i>Aspicarpa hirtella</i>	aspicarpa	Malpighiaceae
	<i>Boerhaavia coccinea</i>	red spiderling	Nyctaginaceae
	<i>Bouchea prismatica</i>	bouchea	Verbenaceae

	SCIENTIFIC NAME	COMMON NAME	FAMILY
FORBS (Cont.)			
	<i>Bouvardia glaberrima</i>	smooth bouvardia	Rubiaceae
	<i>Brickellia</i> spp.	brickellbush	Asteraceae
	<i>Chamaecrista serpens</i> var. <i>wrightii</i>	sensitive pea	Fabaceae
	<i>Cheilanthes fendleri</i>	cloak fern	Pteridaceae
	<i>Cheilanthes</i> spp.	cloak fern	Pteridaceae
	<i>Chenopodium fremontii</i>	lamb's quarter	Chenopodiaceae
	<i>Clitoria mariana</i>	butterfly pea	Fabaceae
	<i>Cnidosculus angustidens</i>	mala mujer	Euphorbiaceae
	<i>Cologania longifolia</i>	narrowleaf tick clover	Fabaceae
	<i>Commelina dianthifolia</i>	western dayflower	Commelinaceae
	<i>Cucurbita digitata</i>	coyote gourd	Cucurbitaceae
	<i>Datura metaloides</i>	sacred datura	Solanaceae
	<i>Eleocharis</i> spp.	spikerush	Cyperaceae
	<i>Eriogonum wrightii</i>	buckwheat	Polygonaceae
	<i>Eryngium heterophylla</i>	button snakeroot	Apiaceae
	<i>Evolvulus alsinoides</i>		Convolvulaceae
	<i>Evolvulus arizonicus</i>	Arizona blue eyes	Convolvulaceae
	<i>Galium wrightii</i>	northern bedstraw	Rubiaceae
	<i>Glandularia gooddingii</i>	verbena	Verbenaceae
	<i>Gnaphalium leucocephalum</i>	white cudweed	Asteraceae
	<i>Gnaphalium wrightii</i>	cudweed	Asteraceae
	<i>Gomphrena</i> sp.	globe amaranth	Amarnathaceae
	<i>Gutierrezia</i> spp.	snakeweed	Asteraceae
	<i>Ipomoea barbatsepala</i>	morning glory	Convolvulaceae
	<i>Ipomoea coccinea</i>	scarlet creeper	Convolvulaceae
	<i>Ipomoea hirsutula</i>	wooly morning glory	Convolvulaceae
	<i>Ipomoea leptotoma</i>	bird's foot morning glory	Convolvulaceae
	<i>Ipomoea longifolia</i>	long leaf morning glory	Convolvulaceae
	<i>Isocoma tenuisecta</i>	burweed	Asteraceae
	<i>Jatropha macrorhiza</i>	Arizona desert potato	Euphorbiaceae
	<i>Kallstroemia grandiflora</i>	Arizona caltrop	Zygophyllaceae
	<i>Krameria parvifolia</i>	range ratany	Krameriaceae
	<i>Machaeranthera</i> spp.	spiny aster	Asteraceae
	<i>Macroptilium gibbosifolium</i>	variableleaf bushbean	Fabaceae
	<i>Milla biflora</i>	Mexican star	Liliaceae
	<i>Oenothera rosea</i>	evening primrose	Onagraceae
	<i>Oxalis albicans</i>	wild oxalis	Oxalidaceae
	<i>Penstemon linarioides</i>	linear leaf penstemmon	Scrophulariaceae
	<i>Phaseolus ritensis</i>	eggleaf stringbean	Fabaceae
	<i>Phaseolus</i> sp.	stringbean	Fabaceae
	<i>Portulaca suffrutescens</i>	portulaca	Portulacaceae
	<i>Portulaca umbraticola</i>	portulaca	Portulacaceae
	<i>Proboscidea</i> sp.	unicorn plant, devil's claw	Pedaliaceae
	<i>Salvia subincisa</i>	sawtooth sage	Lamiaceae

	SCIENTIFIC NAME	COMMON NAME	FAMILY
FORBS (Cont.)			
	<i>Schoenocrambe linearifolia</i>	schoenocrambe	Brassicaceae
	<i>Scirpus</i> sp.	bulrush	Cyperaceae
	<i>Senna covesii</i>	desert senna	Fabaceae
	<i>Senna hirsuta</i>	woolly senna	Fabaceae
	<i>Solanum douglassii</i>	greenspot nightshade	Solanaceae
	<i>Solanum elaeagnifolium</i>	silverleaf nightshade	Solanaceae
	<i>Sphaeralcea</i> spp.	globe mallow	Malvaceae
	<i>Tagetes</i> sp.	marigold	Asteraceae
	<i>Talinum angustissimum</i>	talinum	Portulacaceae
	<i>Talinum aurantiacum</i>	orange fameflower	Portulacaceae
	<i>Tetramerium hispidum</i>	tetramerium	Acanthaceae
	<i>Thalictrum fendleri</i>	Fendler's meadow rue	Ranunculaceae
	<i>Vitis arizonica</i>	Arizona grape	Vitaceae
	<i>Zinnia acerosa</i>	desert zinnia	Asteraceae
TREES & SHRUBS			
	<i>Acacia angustissima</i>	white ball acacia	Fabaceae
	<i>Acacia constricta</i>	whitethorn acacia	Fabaceae
	<i>Acacia greggii</i>	catclaw acacia	Fabaceae
	<i>Aloysia wrightii</i>	oreganillo	Verbenaceae
	<i>Arctostaphylos</i> sp.	manzanita	Ericaceae
	<i>Baccharis salicifolia</i>	seep willow	Asteraceae
	<i>Baccharis sarothroides</i>	desert broom	Asteraceae
	<i>Calliandra eriophylla</i>	fairyduster	Fabaceae
	<i>Celtis pallida</i>	desert hackberry	Ulmaceae
	<i>Celtis reticulata</i>	netleaf hackberry	Ulmaceae
	<i>Chrysothamnus teretifolius</i>	green rabbitbrush	Asteraceae
	<i>Dodonaea viscosa</i>	hopbush	Sapindaceae
	<i>Ericameria laricifolia</i>	turpentine bush	Asteraceae
	<i>Erythrina flabelliformis</i>	coral bean	Fabaceae
	<i>Eysenhardtia orthocarpa</i>	kidney wood	Fabaceae
	<i>Fraxinus velutina</i>	velvet ash; Arizona ash	Oleaceae
	<i>Gossypium thurberi</i>	desert cotton	Malvaceae
	<i>Guardiola platyphylla</i>	Apache plant	Asteraceae
	<i>Hibiscus coulteri</i>	desert rosemallow	Malvaceae
	<i>Indigofera spaerocarpa</i>	Sonoran Indigo	Fabaceae
	<i>Juglans major</i>	Arizona walnut	Juglandaceae
	<i>Juniperus deppeana</i>	alligator juniper	Cupressaceae
	<i>Lasianthaea podocephala</i>	San Pedro daisy	Asteraceae
	<i>Lycium</i> spp.	wolfberry	Solanaceae
	<i>Mimosa biuncifera</i>	catclaw mimosa	Fabaceae
	<i>Mimosa dysocarpa</i>	velvet pod mimosa	Fabaceae

	SCIENTIFIC NAME	COMMON NAME	FAMILY
TREES & SHRUBS (Cont.)			
	<i>Parkinsonia microphylla</i>	yellow paloverde	Fabaceae
	<i>Populus fremontii</i>	Fremont cottonwood	Salicaceae
	<i>Prosopis velutina</i>	velvet mesquite	Fabaceae
	<i>Q. arizonica</i>	Arizona white oak	Fagaceae
	<i>Q. garrya</i>	silktassel	Fagaceae
	<i>Quercus emoryii</i>	Emory oak	Fagaceae
	<i>Rhus aromatica</i>	skunkbush	Anacardiaceae
	<i>Rhus choriophylla</i>	sumac	Anacardiaceae
	<i>Salix exigua</i>	coyote willow	Salicaceae
	<i>Tamarix pentandra</i>	salt cedar	Tamaricaceae
	<i>Ziziphus obtusifolia</i>	graythorn	Rhamnaceae

TEP-Citizen's Interconnect Project

Environmental Training Guidelines Construction Supervisors

- Stay in the designated work areas. Approved work areas, access roads, and staging areas will be clearly marked. All project activities must remain in these areas. Do not work or trespass beyond the signed or fenced restricted work areas.
- Restrict vehicle access to public roadways and designated access roads. Cross-country driving is prohibited.
- No driving or parking within 100 feet of ponds and tanks.
- Do not transfer water from one pond or tank to another or between any other bodies of water.
- No in-stream activity or disposal of construction debris or fill is allowed.
- Store topsoil and trench spoils behind sediment control structures at least 20 feet from any stream bank, including dry washes.
- Check equipment for leaks or heavy surface oil build-up before working in streams or washes.
- The use or transfer of hazardous materials will not be allowed within 100 feet of any stream or wash is prohibited.
- Do not litter. Dispose of trash in designated containers. Uncontained trash can attract wildlife and unwanted pests. Cigarette butts are considered litter, and should be extinguished and disposed of appropriately. All litter and construction debris must be removed from the job site daily.
- No pets or firearms. They are prohibited for job-site protection and protection of wildlife.
- Hunting is prohibited.
- Clearing will be limited to the minimum required to provide a safe construction area. Make sure you know the clearing limit, and if possible, leave plant root systems in place when clearing vegetation.
- It is illegal to harm, harass, pursue, hunt, shoot, wound, trap, kill capture, or collect wildlife officially listed as threatened or endangered. Violation of threatened and endangered special laws can result in penalties of up to \$100,000 and/or one year in jail.
- Do not approach or feed wildlife. Keep away from their burrows and nests. Do not harm or kill any wildlife encountered.
- If animal is harmed or found harmed, contact your Construction Supervisor or the Environmental Inspector. Do not attempt to move the animal yourself.

APPENDIX D

APPENDIX D. Federally Listed, Proposed, and Candidate Species under jurisdiction of the U.S. Fish and Wildlife Service in Pima County, Arizona as of 14 August 2002, excluded from further consideration.

COMMON NAME	SCIENTIFIC NAME	STATUS	HABITAT	JUSTIFICATION
PLANTS				
Canelo Hills ladies' tresses	<i>Spiranthes delitescens</i>	Endangered	Finely grained, highly organic, saturated soils of cienegas. Potential habitat occurs in Sonora, Mexico.	No habitat present.
Huachuca water umbel	<i>Lilaeopsis schaffneriana</i> ssp. <i>recurva</i>	Endangered	An emergent aquatic plant that requires marshy wetlands.	No habitat present.
Kearney's blue star	<i>Amsonia kearneyana</i>	Endangered	Known only from the Baboquivari Mountains.	ROW is outside of known range.
Nichol's Turk's head cactus	<i>Echinocactus horizonthalonius</i> var. <i>nicholii</i>	Endangered	Dependent on limestone substrates in desert hills.	No habitat present.
FISH				
Desert pupfish	<i>Cyprinodon macularius</i>	Endangered	Shallow springs, small streams, and marshes. Tolerates saline and warm water.	No habitat present in area.
Gila chub	<i>Gila intermedia</i>	Proposed Endangered	Small streams and cienegas; prefer deeper pools with cover.	No habitat present in area.
Loach minnow	<i>Tiaroga cobitis</i>	Threatened	Requires perennial streams with swift water over cobble or gravel	No habitat present in area.
Sonoran Chub	<i>Gila ditaenia</i>	Threatened	Most commonly found in deep, permanent pools with bedrock-sand substrates and free of floating algae.	In U.S, limited to Sycamore Canyon and its tributaries.
Spikedace	<i>Meda fulgida</i>	Threatened	Requires perennial streams with swift velocities over sand and gravel.	No habitat present in area.
AMPHIBIANS				
Sonoran tiger salamander	<i>Ambystoma tigrinum stebbinsi</i>	Endangered	Stock tanks and impounded cienegas in San Rafael Valley, Huachuca Mountains at 4000-6300 ft.	ROW is outside of known range. This species is not known to occur in the Nogales RD.
Chiricahua leopard frog	<i>Rana chiricahuensis</i>	Threatened	Perennial pools, springs, stock tanks and ponds above 3,500' elevation.	No occupied habitat within ROW and no reintroductions planned.

APPENDIX D. Federally Listed, Proposed, and Candidate Species under jurisdiction of the U.S. Fish and Wildlife Service in Pima County, Arizona as of 14 August 2002, excluded from further consideration.

COMMON NAME	SCIENTIFIC NAME	STATUS	HABITAT	JUSTIFICATION
BIRDS				
Bald eagle	<i>Haliaeetus leucocephalus</i>	Threatened	Large trees or cliffs near water (reservoirs, rivers, and streams) with abundant prey.	Winter surveys of Peña Blanca and Arivaca Lakes were conducted in 1994-1996, 1998, 2000-2002. No bald eagles have been observed.
California brown pelican	<i>Pelecanus occidentalis californicus</i>	Endangered	Coastal land and islands; species is found around many Arizona lakes and rivers.	No habitat present in area.
Masked bobwhite	<i>Colinus virginianus ridgewayi</i>	Endangered	Only known Arizona population has been reintroduced on Buenos Aires Natl. Wildl. Refuge	ROW is outside of known range.
Mexican spotted owl	<i>Strix occidentalis lucida</i>	Threatened	Montane forests and woodlands and canyon bottoms generally above 4,000-ft. elevation	No habitat within 1 mile of proposed ROW.
Mountain plover	<i>Charadrius montanus</i>	Proposed Threatened	Open arid plains, short grass prairies, and cultivated farms.	No habitat present in area.
Northern apolomado falcon	<i>Falco femoralis septentrionalis</i>	Endangered	Grassland and savannah habitats	No recent confirmed reports for Arizona.
MAMMALS				
Ocelot	<i>Felis pardalis</i>	Endangered	Prefers humid tropical & sub-tropical habitats; typically found at higher elevations.	ROW is outside of known range.
Jaguarundi	<i>Felis yagouaroundi tolteca</i>	Endangered	Deciduous forests, riparian areas, swampy grasslands, upland drysavannahs, etc.	ROW is outside of known range.
Sonoran pronghorn	<i>Antilocapra americana sonoriensis</i>	Endangered	Grassy desertscrub in northwestern Sonora and adjacent Arizona borderlands, mainly Yuma Co.	ROW is outside of known range.

STATUS DEFINITIONS: ENDANGERED SPECIES ACT

Endangered: Imminent jeopardy of extinction.

Threatened: Imminent jeopardy of becoming endangered.

Proposed: Proposed Rule has been published in Federal Register to list as Threatened or Endangered.